

Impacts of immunostimulant yeast (*Saccharomyces cerevisiae*) supplemented feed on growth and blood profile of Java Barb

by Tita Elfitasari

Submission date: 13-Jun-2022 10:15AM (UTC+0700)

Submission ID: 1855711418

File name: 2021_-_Diana_R_-_Pak_Roy_-_Tita.pdf (223.57K)

Word count: 4579

Character count: 23088

Impacts of Immunostimulant Yeast (*Saccharomyces cerevisiae*) Supplemented Feed on Growth and Blood Profile of Java Barb (*Barbonymus gonionotus*)

Diana Rachmawati^{1,*}, Roy Hendroko Setyobudi², Juris Burlakovs³, Tita Elfitasari¹ and Agus Heri Purnomo⁴

¹Department of Aquaculture, Faculty of Fisheries and Marine Sciences, Diponegoro University, Jl. Prof. Soedarto SH, Tembalang, Semarang 50275, Indonesia; ²Department of Agriculture Science, Postgraduate Program, University of Muhammadiyah Malang, Jl. Raya Tlogomas No 246, Malang 65114, Indonesia; ³Department of Water Management, Estonian University of Life Sciences, Tartu, Estonia; Friedrich Reinhold Kreutzwald 1a, 51014 Tartu, Estonia; ⁴Ministry of Marine Affairs and Fisheries Republic of Indonesia, The Research Center for Marine and Fisheries Socio-economics, Mina Bahari 1-Building, Jl. Pasir Putih 1, Ancol, Jakarta 14430, Indonesia

Received: Feb 20, 2021; Revised: April 29, 2020; Accepted May 1, 2021

Abstract

12

The research aims to analyze impacts of immunostimulant yeast (*Saccharomyces cerevisiae*) supplemented feed on growth rate, blood profile in Java Barb [*Barbonymus gonionotus* (Bleeker, 1850)] fingerlings. The Java Barb of 200 fingerlings have an average weight of $5.64 \text{ g} \pm 0.87 \text{ g}$, and an average length of $6.32 \text{ cm} \pm 0.58 \text{ cm}$. The experiments employed a Completely Randomized Design with five treatments, and each had four repetitions for a period of 49 d (days). Various dosages of *S. cerevisiae* were added to the feed. The treatments were T1 (0 g kg⁻¹ feed), T2 (1 g kg⁻¹ feed), T3 (2 g kg⁻¹ feed), T4 (3 g kg⁻¹ feed), and T5 (4 g kg⁻¹ feed). The results showed that yeast as supplemented feed could improve the growth and blood profiles of Java Barb. The blood profiles include improvement in total blood cells (TCC μL^{-1}), red blood cells (RBC), white blood cells (WBC), hematocrit (HTC) compared to control. The highest value of efficiency of feed utilization (EFU) was 72.36 % and followed by FCR (feed conversion ratio), PER (protein efficiency ratio), RGR (relative growth rate), and SR (survival rate) with the values of (1.63, 4.21, 4.18) % d⁻¹, and 92.33 % respectively. The optimal dosages of *S. cerevisiae* in the feed for EFU, FCR, PER, and RGR ranged from 2.38 g kg⁻¹ feed to 3 g kg⁻¹ feed.

Keywords: Digestibility, Enzyme, Fisheries feed, Immunity, Resistance

1. Introduction

Java Barb [*Barbonymus gonionotus* (Bleeker, 1850)] is an Indonesian native fish that is easily cultivated; hence, farmers have practiced intensively in aquaculture (Rachmawati *et al.*, 2019b). One of the problems faced by the aquaculture farmers was decreased in water quality due to unconsumed fish feeds and water waste. This condition could inhibit fish growth and rising fish diseases that could decrease fish production. Thus, it made farmers lose in profit. The studies to administrate the fish growth have been done through giving biofloc, probiotic, herb, and immunostimulant. Biofloc was implemented on fish and shrimp (Kuhn *et al.*, 2010). Moreover, de La Banda *et al.* (2010) has applied probiotic on fish and shrimp cultivation. Furthermore, Goda *et al.* (2012) also reported the *S. cerevisiae* supplemented feed could boost fish appetite; therefore, it improved growth and survival rate.

To spur growth and immunity in the fish, fish farmers could utilize immunostimulants that can be obtained from seaweed, bacteria, or yeast. Manoppo and Magdalena (2015) also discovered that the utilization of immunostimulants could raise growth and the immune system in the fish and crustaceae. According to Abu-Elala

et al. (2013), *S. cerevisiae* is one of the immunestimulants. In addition, Tewary and Patra (2011) also suggested that *S. cerevisiae* can trigger digestibility due to digestive enzymes to boost fish growth. The *S. cerevisiae*-supplemented feed could boost the digestibility of feed and protein. It could also increase efficiency utilization of the feed and the growth (Razak *et al.*, 2017).

The immune response rate of fish could be increased by adding *S. cerevisiae* in the feed, as reported by Abu-Elala *et al.* (2013). Next, the *S. cerevisiae* could be used as an immunostimulant, because it is rich in elements such as β -1-3 glucan (50 % to 60 %). Moreover, it could generate its immunity in the fish and crustacean in which β -glucan in the yeast was able to boost the immunity and disease resistance of the fish (Manoppo and Magdalena, 2015). Moreover, Sheikhzadeh *et al.* (2012) mentioned that *S. cerevisiae* contains immunostimulants such as β -1,3 glucan, nucleic acid, manna oligosaccharide, chitin, non-starch nucleic acid, and polysaccharide. The β -1,3 glucan effectively increased the immune system in some species of fish at the dosage of 1 g kg⁻¹ feed (Dhanaraj *et al.*, 2010). Jarmolowicz *et al.* (2011) disclosed that the addition of beer yeast at the dosage of 4 % to 6 % could boost non-specific immune system in the fingerlings of pikeperch [*Sander lucioperca* Linnaeus, 1758)].

* Corresponding author e-mail: dianarachmawati964@gmail.com

Some researches on the *S. cerevisiae*-supplemented feed to boost growth and immunity were disclosed by Tewary and Patra (2011) in *Labeo rohita*, F. Hamilton, 1822; Tawwab *et al.* (2010) in *Sarotherodon galileaus*, Linnaeus, 1758; Manoppo and Kolopita (2016) in *Cyprinus carpio*, Linnaeus, 1758; Jarmolowicz *et al.* (2011) in *Sander lucioperca* Linnaeus, 1758; Azevedo *et al.* (2016) in *Oreochromis niloticus*, Linnaeus 1758; Rachmawati *et al.* (2019a) in *Pangasius hypophthalmus* Sauvage, 1878; and Rachmawati *et al.* (2019b) in *Barbomyrus gonionotus* Bleeker, 1850. However, previous research only observed the growth of cultivated fish, and research on the effect of blood profiles has never been carried out. Therefore, it is important to do this research which studies the impact of yeast on the blood profile of Java Barb.

Some benefits of using *S. cerevisiae* as an immunostimulant in aquaculture are that it does not leave any residues in fish and the environment, and hazard free for human beings as well. Therefore, the utilization of *S. cerevisiae* as an immunostimulant to increase growth is very important. This study aims to analyze the impacts of *S. cerevisiae* supplemented feed on growth rate and blood profile in Java Barb fingerlings.

24

2. Materials and Method

The study was conducted at the Laboratory for Fish Health and Environment Assessment, Muntulan, Central Java, Indonesia from April until June 2019. This experiment utilized 500 sample fingerlings of Java barb with an average weight of $5.64 \text{ g} \pm 0.87 \text{ g}$, and an average length of $6.32 \text{ cm} \pm 0.58 \text{ cm}$. (Rachmawati *et al.*, 2017). To adjust to feeding and a new environment, the sample fish was first acclimated for 1 wk (week). During acclimatization, fish were fed with commercial feed using ad satiation method. To maintain the water quality, water was being siphoned before feeding. Before the study was conducted, the sample fish fasted for 1 d (day) to clean the metabolism residues hence the initial weight was not affected by the waste weight. The yeast materials for treatments were commercial yeast (*S. cerevisiae*) brand Saf Instant produced by Saf Indonusa - Lessafre Global Group, Indonesia.

Completely Randomized Designed (CRD) was implemented in this research. There were five treatments with four replications. The treatments were a supplementation of yeast (*S. cerevisiae*) into the feed. The feeds used were commercial with brand name Comfeed fish feed, produced by JAPFA Comfeed Indonesia Tbk., composed of 30 % of raw protein, 2 % fat, 3 % raw fiber, 13 % ash, and 12 % water. Treatments were included: T₁ (0 g kg⁻¹ feed), T₂ (1 g kg⁻¹ feed), T₃ (2 g kg⁻¹ feed), T₄ (3 g kg⁻¹ feed), and T₅ (4 g kg⁻¹ feed). The dosages of the *S. cerevisiae* in this study were modified from Rachmawati *et al.* (2019a) study. The study recorded that 1 g kg⁻¹ feed supplementation of *S. Cerevisiae* was the best dosage to produce the highest feed usage efficiency in fingerling of catfish (*Pangasius hypophthalmus* Sauvage, 1878).

After the yeast was added, the protein analysis per treatment was T₀ (30 %), T₁ (30.37 %), T₂ (30.46 %), T₃ (30.39 %), and T₄ (30.53 %). The preparation of the feed was by weighing the *S. cerevisiae* based on the treatments. The weighed yeast was diluted with pure

water. The 100 mL pure water was diluted in 1 kg feed (Manurung *et al.*, 2013). The diluted yeast was evenly sprayed into the feed based on the Rachmawati *et al.*, (2019b) method. Firstly, the feed was put on the plastic tray and then sprayed by suspense yeast. When feed was being sprayed with yeast, the tray was shaken and the feed and the yeast were mixed. Then, the mixture was dried up by letting it at room temperature of 28 °C to 30 °C. After that, the mixture was put in a plastic bag. Then, the bag was labeled by treatment and stored in the refrigerator until it was ready to use. The fix feeding method was used to feed the fingerlings. The method was based on 5 % of the fish weight. The three times daily feeding implemented were in the morning (7 am), afternoon (2 pm), and evening (6 pm). The scale of the fish weight was carried out every week for 49 d. Sampling was carried out once 1 wk; therefore, a 7 wk observation is required to obtain more accurate data.

The containers used were made from fiberglass by dimension 1.5 m × 1.5 m × 1 m. The fish was reared in 20 containers with a stock density of 25 fish m⁻³. Freshwater was utilized as cultivation media. Before the water was used, it has been deposited for several days in a reservoir.

The observation on blood profile has been conducted at the beginning (1st d) and the end (49th d) of the study. The observed parameters included total cells (TCC μL^{-1}), red blood cells (RBC), white blood cells (WBC), hematocrit (HCT) (referring to Mohammed *et al.*, 2013 on blood profile methodology).

The observed variables included feed efficiency (EF), the ratio of feed conversion (FCR), the protein efficiency ratio (PER), relative growth rate (RGR), and survival rate (SR) which was measured every week. Those variable observations were referring to Rachmawati *et al.* (2018) and Rachmawati *et al.* (2019b), while analysis of blood profile and water quality consisting of temperature, pH, and dissolved oxygen was measured every morning and evening, whereas ammoniac was measured in 1st d, 28th d and 49th d based on the method of APHA (1992). Water quality was maintained through the daily siphon and 30 % of water change every week after sampling. The measure variables are EFU (efficiency of feed utilization) in Equation (1) and RGR (relative growth rate) in Equation (2):

$$EFU = 100 \left(\frac{W_2 - W_1}{QF} \right) \quad (1)$$

$$RGR = 100 \left(\frac{W_2 - W_1}{T \times W_1} \right) \quad (2)$$

Note: W₁ and W₂ are the initial and final weight, respectively. QF is the feed amount consumed, and T is the days during the feeding period. FCR (feed conversion ratio), PER (protein efficiency ratio), SR (survival rate) are calculated by Equation (3), Equation (4), and Equation (5).

$$FCR = 100 \left(\frac{F_1}{WG} \right) \quad (3)$$

Note: FI is feed intake (g) and WG is weight gain (g);

$$PER = 100 \left(\frac{W}{PI} \right) \quad (4)$$

Note: WG is weight gain (g) and PI is protein intake (g);

$$SR = 100 \left(\frac{C_2}{C_1} \right) \quad (5)$$

29

¹⁰: C₁ is initial count of fish and C₂ final count of fish.

Analysis of variance (ANOVA) and Duncan's Multiple Range Test were used to analyze the observed variables (Rachmawati *et al.*, 2019b). The polynomial orthogonal test using SAS9 and Maple12 software was used to calculate the optimal dosage of immunostimulant yeast (*S. cerevisiae*). Water quality parameters were descriptively explained by comparing the rearing conditions to determine the viability.

3. Results and Discussions

Table 1. displayed the observed variables of EFU, FCR, PER, RGR, and SR in treatment T₀, T₁, T₂, T₃, and T₄. The polynomial orthogonal test results of *S. cerevisiae*-¹⁵ implemented feed on EFU, FCR, PER, and RGR were displayed ¹⁹, Figure 1, Figure 2, Figure 3, and Figure 4. Figure 1 showed that the optimal dosage of *S. cerevisiae* in the feed for EFU was 2.38 g kg⁻¹ feed with the EFU value as much as 67.67 %. The best value of FCR (1.58) was obtained from *S. cerevisiae* with 2.43 g kg⁻¹ feed, as shown in Figure 2.

Table 1 shows that adding yeast to feed has a significant effect on EFU, FCR, PER, and RGR, but does not show an effect on SR.

Figure 3 displayed the relationship between the dosage of yeast and the values of PER. The optimal dosage of *S. cerevisiae* for PER was 2.62 g kg⁻¹ feed giving the value of 4.08. Based on Figure 4, one could calculate the optimal dosage of *S. cerevisiae* for RGR. The optimum dosage was 3 g kg⁻¹ feed giving the value of 4.18 % d⁻¹.

Table 1. The values of the variables

Experiment	Treatments				
Data	T ₀	T ₁	T ₂	T ₃	T ₄
EFU (%)	48.24 ± 0.18 ^d	59.75 ± 0.27 ^c	63.35 ± 0.31 ^b	72.36 ± 0.19 ^a	55.63 ± 0.15 ^c
FCR	2.53 ± 0.32 ^c	2.18 ± 0.25 ^b	2.01 ± 0.25 ^b	1.63 ± 0.18 ^a	2.12 ± 0.27 ^b
PER	1.96 ± 0.28 ^d	2.85 ± 0.24 ^c	3.27 ± 0.23 ^b	4.21 ± 0.27 ^a	2.59 ± 0.25 ^c
RGR (% d ⁻¹)	1.67 ± 0.14 ^d	2.09 ± 0.15 ^c	3.52 ± 0.22 ^b	4.18 ± 0.25 ^a	3.48 ± 0.22 ^b
SR (%)	75.33 ± 3.52 ^a	90.33 ± 2.67 ^a	90.33 ± 2.72 ^a	92.33 ± 2.78 ^a	90.33 ± 2.43 ^a

Note: The mean values with a different superscript in the same column showed significant difference (P < 0.05)




Figure 1. The relation between the immunostimulant yeast (*S. cerevisiae*) and EFU in the Java Barb fingerlings




Figure 2. The relation between the immunostimulant yeast (*S. cerevisiae*) and FCR in the Java Barb fingerlings




Figure 3. The relation between the immunostimulant yeast (*S. cerevisiae*) and PER in the Java Barb fingerlings




Figure 4. The Relations between the immunostimulant yeast (*S. cerevisiae*) and RGR in the Java Barb fingerlings

The enrichment of *S. cerevisiae* with the dosages of 8 g kg^{-1} feed to 4 g kg^{-1} feed (T1, T2, T3, T4) in the Java barb fingerlings had higher values of EFU (55.63 % to 72.36 %) than those given the zero-supplementation yeast (48.24 %, as in the treatment T0). It was noted that the *S. cerevisiae* supplemented feed could raise the efficiency of feed utilization. Razak *et al.* (2017) discovered that the *S. cerevisiae* could boost enzymatic activities in the digestive track. Hence, it enhanced the decomposition of complex nutrients into simpler nutrients as proclaimed by Tewary and Patra (2011). The simpler nutrients were easier to absorb, and in turn this enhanced the efficiency of feed utilization. The dosage of 3 g kg^{-1} feed (T3) produced the highest EFU (72.36 %), followed by the treatments T2, T1, T4, and T0 that had values of 63.35 %, 59.75 %, 55.63 % and 48.24 %, respectively. The optimal dosage of yeast in the feed that resulted in the highest EFU may cause the dosage appropriate to enhancing enzyme activities in the digestive track and then improve the efficiency of feed usage. Welker *et al.* (2012) discovered

that *S. cerevisiae* as an immunostimulant could enhance the production of the enzyme in the digestive tract to boost digestibility and absorption for nutrients, amino acids, vitamins, and enzymes. Similar results from other studies were reported by Tawwab *et al.* (2010) in *Sarotherodon galileus* Linnaeus 1758, Tewary and Patra (2011) in *Labeo rohita* F. Hamilton 1922, and Rachmawati *et al.* (2019a) in *Pangasius hypothalamus* Sauvage 1978. Results of previous experiments showed that the addition of yeast to feed increases EFU.

The *S. cerevisiae* supplemented feed decreased the FCR of Java barb fingerlings. The treatment T3 with the dosage of 3 g kg^{-1} feed generated the lowest FCR. The dosage was to make maximum feed usage efficiency, so it decreased FCR. Razak *et al.* (2017) supported this discovery. This study found that the *S. cerevisiae*-supplemented feed could raise the efficiency of feed usage and decreased the ratio of feed conversion. In addition, Jarmolowicz *et al.* (2011) discovered that the *S. cerevisiae* supplemented feed caused FCR to increase due to the increase of protein digestibility. Essa *et al.* (2010) also disclosed that the *S. cerevisiae*-supplemented feed raised the FCR and the efficiency of feed utilization.

The protein efficiency ratio (PER) is a number calling the total weight of fish produced per unit of protein in the feed (Tiamiy *et al.*, 2014). Various dosages of *S. cerevisiae* of 1 g kg^{-1} feed to 4 g kg^{-1} feed given to Java barb fingerlings generated higher PER (2.59 to 4.21) compared to the PER (1.96) without the addition of yeast. Goda *et al.* (2012) mentioned that the *S. cerevisiae*-supplemented feed could surge protein digestibility; therefore, it increased the ratio of protein efficiency. Moreover, Rachmawati *et al.* (2019b) stated that the *S. cerevisiae*-supplemented feed could enhance the efficiency of feed utilization and the ratio of protein efficiency. The highest PER (4.21) was achieved in T3 (3 g kg^{-1} feed), while the lowest PER (1.96) was attained in T0 (0 g kg^{-1} feed). The highest PER was because of the right dosage of the yeast to generate the ratio of protein efficiency, while the lowest PER was due to the absence of yeast.

The results on the RGR of the supplementation of *S. cerevisiae* ranged from 1 g kg^{-1} feed to 4 g kg^{-1} feed and between $2.09 \% \text{ d}^{-1}$ and $4.18 \% \text{ d}^{-1}$. These values were higher than without yeast supplementation by $1.67 \% \text{ d}^{-1}$. The *S. cerevisiae* supplemented feed could raise RGR, which was due to the increases in protein digestibility and efficiency of feed utilization (Manoppo and Magdalena, 2015). Moreover, Rajagukguk *et al.*, (2017) disclosed that the existence of yeast (*S. cerevisiae*) in the fish digestive system could boost enzymatic activities. It can hike protein digestibility and efficiency of feed usage, and in turn, it raised RGR. The treatment T3 (3 g kg^{-1} feed) generating the highest RGR was suggested due to the effective dosage of the yeast (*S. cerevisiae*).

The *S. cerevisiae* supplemented feed was not significant ($P > 0.05$) on the SR of Java barb fingerlings. It was suggested that the feed was not a factor for SR. The factor that affected SR was abiotic one, such as the ability to adapt to the environment, handling, density, competitors, diseases, ages, and the existence of the predators (Abu-Elala *et al.* (2013). The SR of Java barb fingerlings that were fed by supplementation of *S. cerevisiae* ranged from 90.33 % to 92.33 %, while the survival rate for those that were not fed with the

supplementation of *S. cerevisiae* was 73.33 % (Table 1). It was suggested that the yeast (*S. cerevisiae*) contained β-glucan as an agent of immunostimulant to increase the immune system; therefore, the SR was high. This phenomenon supported by Manoppo and Magdalena (2015) that the enrichment of β-glucan was able to be an immunostimulant to improve the immune system in the fish. The results of the blood profile were higher after the fingerlings were fed with the enrichment of *S. cerevisiae* as in Table 2.

The results of the blood profile measurement consisted of total cell count (TCC/ μL^{-1}), red blood cells (RBC), white blood cells (WBC), and hematocrit (HCT). Those were displayed in Table 2.

Table 2. Total cell count (TCC/ μL^{-1}), red blood cells (RBC), and white blood cells (WBC), hematocrit (HCT) in the Java Barb fingerlings

Treatments (g kg $^{-1}$ feed)	TCC/ μL ($\times 10^6$)	RBC ($\times 10^6$)	WBC ($\times 10^5$)	HCT (%)
T ₁ (0)	1.39 ^b	1.92 ^b	1.38 ^b	24.49 ^b
T ₂ (1)	3.23 ^a	3.25 ^a	2.63 ^a	35.13 ^a
T ₃ (2)	3.45 ^a	3.37 ^a	2.37 ^a	36.74 ^a
T ₄ (3)	3.65 ^a	3.39 ^a	2.65 ^a	37.98 ^a
T ₅ (4)	3.56 ^a	3.29 ^a	2.83 ^a	35.83 ^a
±SD	0.127	0.132	0.263	0.221

Note: The mean values with a different superscript in the same column showed significant difference ($P < 0.05$)

Table 2 showed that the fingerlings that were fed with the *S. cerevisiae* supplemented feed had a significant increase in the total cell count (TCC/ μL^{-1}), the red blood cells (RBC), the white blood cells (WBC), and the hematocrit (HCT). It was revealed that the *S. cerevisiae*-supplemented feed increased non-specific immune response in the Java barb fingerlings. The results show that *S. Cerevisiae* is rich in β-1-3 glucan (50 % to 60 %) that increased fish immune system; therefore, fish resisted the pathogen. It was proven by the value of SR that was higher than without *S. Cerevisiae* supplementation. Manoppo and Magdalena (2015) reported that the *S. cerevisiae*-supplemented feed increased non-specific immune response. Abu-Elala *et al.* (2013) also reported that the Java barb fed with *S. cerevisiae* supplemented feed increased erythrocyte, hemoglobin, and leukocyte. Moreover, Welker *et al.* (2012) discovered that the catfish fed with *S. cerevisiae*-supplemented feed for 1 wk had higher red and white blood cells than those with no *S. cerevisiae* supplemented feed.

The observation of water quality during the experimental period was in an infeasible condition for Java Barb cultivation as indicated by the standard quality in the literature. The suitable condition of the water quality affected by the water quality was controlled according to the requirement of Java Barb need.

4. Conclusion

The supplementation of immunostimulant yeast (*S. cerevisiae*) in the feed could raise the growth and blood profiles such as total cell count, red blood cells, and white blood cells in the Java barb fingerlings. The optimal

dosages of *S. cerevisiae* in feed for EFU, FCR, PER, and RGR ranged from 2.38 g kg $^{-1}$ feed to 3 g kg $^{-1}$ feed.

Acknowledgment

Appreciation is sincerely conveyed to the Head of Laboratory for Fish Health and Environment Assessment, Muntilan, Central Java, Indonesia who has provided the facility to carry out this research.


References

- Abu-Elala, N, Marzouk M and Moustafa M. 2013. Use of different *Saccharomyces cerevisiae* biotic forms as immunomodulator and growth promoters for *Oreochromis niloticus* challenged with some fish pathogens. *J Vet Med Sci.* 1:21-29.
- Azevedo RV, Filho JCF, Pereira SL, Cardoso LD, Andrade DR and Júnior MVV. 2016. Dietary mannan oligosaccharide and *Bacillus subtilis* in diets for Nile tilapia (*Oreochromis niloticus*). *Acta Sci: Veterinariae.* 38(4):347-353.
- De La Banda IG, Lobo C, Leon-Robio JM, Tapia-Paniagua S, Balebona MC, Morinigo, MA, Moreno-Venta X, Lukas LM, Linares F, Arce F and Ariji S. 2010. Influence of two closely related probiotics on juvenile Senegalese sole (*Solea senegalensis*, Kaup 1858) performance and protection against *Photobacterium damsela* ssp. *Piscicida*. *Aquac.* 306:281-288.
- Dhanaraj M, Haniffaa MA, Singha SVA, Arockiarajb AJ, Ramakrishnanan CM, Seetharamana S and Arthimanjua R. 2010. Effect of probiotics on growth performance of koi carp (*Cyprinus carpio*). *J App Aquac.* 22(3): 202-209.
- Essa MA, El-Serafy SS, El-Ezabi MM, Daboor SM, Esmael NA and Lali SP. 2010. Effect of different dietary probiotics on growth, feed utilization, and digestive enzyme activities of Nile tilapia, *Oreochromis niloticus*. *J World Aquac Soc.* 5(2): 143-162.
- Goda AMA, Mabrouk HAH, Wafa MAE and El-Afifi TM. 2012. Effect of using baker's yeast and exogenous digestive enzymes as growth promoters on growth, feed utilization, and hematological indices of Nile tilapia. *Oreochromis niloticus* Fingerling. *J Agric Sci Technol B.* (2):15-28.
- Jarmolowicz S, Zakes Z, Siwicki A, Kowalska A, Hopko M, Glabski E, Demska Z and Partyka K. 2012. Effects of brewer's yeast extract on growth performance and health of juvenile pikeperch *Sander lucioperca* (L.). *Aquac Nutr.* 18(4): 457-464
- Kuhn DD, Lawrence AL, Boardman GD, Patnaik S, Marsh L and Flick Jr. GJ. 2010. Evaluation of two types of bio floc derived from biological treatment of fish effluent as feed ingredients for Pacific white shrimp *Litopenaeus vannamei*. *Aquac.* 303:28-33.
- Manoppo H and Magdalena EF. 2015. Composting bread yeast in feed increases nonspecific immune response and Tilapia growth. *Jurnal Veteriner.* 16(2):204-211.
- Manoppo H and Kolopita MEF. 2016. Use of bread yeast (*Saccharomyces cerevisiae*) as immunostimulant to improve the resistance of carp (*Cyprinus carpio* L) against bacterial infection *Aeromonas hydrophila*. *Jurnal Budidaya Perairan.* 4(3):37-47.
- Manurung UN, Manoppo H and Tumbol RA. 2013. The use of baker's yeast (*Saccharomyces cerevisiae*) in enhancing non specific immune response and growth of nile tilapia (*Oreochromis niloticus*). *Jurnal Budidaya Perairan.* 1(1):8-14.
- Mohammed H, Olivares-Fuster O, Lafrentz S and Arias CR. 2013. New attenuated vaccine against columnaris disease in fish: Choosing the right parental strain is critical for vaccine efficacy. *Vaccine.* 31:5276-5280.
- Rachmawati D, Istiyanto S and Maizirwan Mel. 2017. Effect of phytase on growth performance, feed utilization efficiency and

- nutrient digestibility in fingerlings of *Chanos chanos* (Forsskal 1775). *Philipp J Sci.* **146** (3): 237-245.
- Rachmawati D, Prihanto AA, Roy HS and Anne O. 2018. Effect of papain enzyme in feed on digestibility of feed, growth performance, and survival rate in post larvae of freshwater Lobster [*Cherax quadricarinatus* (Von Martens, 1868)]. *Proc Pakistan Acad Scies B.* **55** (3): 31-39
- Rachmawati D, Istiyanto S, Ristiawan AN and Susilowati T. 2019a. Effects of *Saccharomyces cereviceae* incorporated diet on growth performance, apparent digestibility coefficient of protein and survival rate of catfish (*Pangasius hypothalamus*). *Aquacultura Indonesiana.* **20** (1):8-14.
- Rachmawati D, Hutabarat H, Samidjan I, Herawati VE and Windarto S. 2019b. The effects of *Saccharomyces cerevisiae*-enriched diet on feed usage efficiency, growth performance and survival rate in Java barb (*Barbonyx gonionotus*) fingerlings. *Aquac Aquar Conserv Legis.* **12** (5):1841-1849.
- Rajagukguk BB, CLumenta C and Mokolensang JF. 2017. Utilization of yeast (*Saccharomyces cerevisiae*) in feed formulations in increasing the growth of tilapia (*Oreochromis niloticus*). *Jurnal Budidaya Perairan.* **5**(3):44-49.
- Razak AP, Kreckhoff RL and Watung JC. 2017. Administration of oral immunostimulant bread yeast (*Saccharomyces cerevisiae*) to increase goldfish growth (*Cyprinus carpio* L.). *Jurnal Budidaya Perairan.* **5**(2):27-36.
- Sheikhzadeh AE, Heidarich M, Pashaki AK, Nofouzi K, Farshabi MA and Akbari M. 2012. Hilyses fermented *Saccharomyces cereviceae*, enhances the growth performance and skin non-specific immune parameters in rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol.* **32** (6):1083-1087.
- Tawwab MA, Mousa MAA and Mohammed MA. 2010. Use of live baker's yeast, *Saccharomyces cerevisiae*, in practical diet, to enhance the growth performance of galilee tilapia, *Sarotherodon galilaeus* (L.), and its resistance to environmental copper toxicity. *J World Aquac Soc.* **41**(S2):214-223.
- Tewary A and Patra BC. 2011. Oral administration of baker's yeast (*Saccharomyces cerevisiae*) acts as a growth promoter and immunomodulator in *Labeo rohita* (Ham.). *J Aquac Res Dev.* **2**(1):1-7.
- Tiamiyu, LO, Victoria OA, Victor TO and Saidu U. 2014. Effect of various levels of raw *Citrullus lanatus* seed meal diets on growth performance of *Cyprinus carpio* Fingerlings. *Jordan J Biol. Sci.* **7**(4):269-274.
- Welker T L, Lim C, Yildirim-Aksoy M and Klesius PH. 2012. Effect of short-term feeding duration of diets containing commercial whole-cell yeast or yeast subcomponents on immune function and disease resistance in channel catfish, *Ictalurus punctatus*. *J Anim Physiol Anim Nutr (Berl).* **96**:159-171.

Impacts of immunostimulant yeast (*Saccharomyces cerevisiae*) supplemented feed on growth and blood profile of Java Barb

ORIGINALITY REPORT



PRIMARY SOURCES

- | | | |
|---|---|-----|
| 1 | www.jurnal.unikal.ac.id
Internet Source | 1 % |
| 2 | Submitted to Padjadjaran University
Student Paper | 1 % |
| 3 | ijasr.um.ac.ir
Internet Source | 1 % |
| 4 | Submitted to Kenyatta University
Student Paper | 1 % |
| 5 | www.ifrj.upm.edu.my
Internet Source | 1 % |
| 6 | lrrd.org
Internet Source | 1 % |
| 7 | sphinxsai.com
Internet Source | 1 % |
| 8 | Ganqiu Lan, Norhani Abdullah, Syed Jalaludin, Yin Wan Ho. " Effects of freeze-dried culture and Natuphos phytase supplementation on the performance and nutrient utilisation of | 1 % |

broiler chickens ", Journal of the Science of Food and Agriculture, 2012

Publication

-
- 9 L Setijaningsih, E Setiadi, I Taufik, Mulyasari. "The effect of garlic Allium sativum addition in feed to the growth performance and immune response of tilapia Oreochromis niloticus", IOP Conference Series: Earth and Environmental Science, 2021
Publication <1 %
- 10 koreascience.or.kr <1 %
Internet Source
- 11 submission.intelaquares.com <1 %
Internet Source
- 12 www.pertanika.upm.edu.my <1 %
Internet Source
- 13 citeseerx.ist.psu.edu <1 %
Internet Source
- 14 onlinelibrary.wiley.com <1 %
Internet Source
- 15 www.mdpi.com <1 %
Internet Source
- 16 www.researchgate.net <1 %
Internet Source
- 17 eprints.nottingham.ac.uk <1 %
Internet Source

-
- 18 Aditya P. Yanuar, Henky Manoppo. "Respon kebal nonspesifik ikan mas yang diberi imunostimulan ragi roti secara oral", e-Journal BUDIDAYA PERAIRAN, 2017 <1 %
Publication
-
- 19 Mohsen Abdel-Tawwab. "Interactive effects of dietary protein and live bakery yeast, *Saccharomyces cerevisiae* on growth performance of Nile tilapia, *Oreochromis niloticus* (L.) fry and their challenge against *Aeromonas hydrophila* infection", Aquaculture International, 2011 <1 %
Publication
-
- 20 R. L. Walzem, C. J. Dillard, J. B. German. "Whey Components: Millennia of Evolution Create Functionalities for Mammalian Nutrition: What We Know and What We May Be Overlooking", Critical Reviews in Food Science and Nutrition, 2002 <1 %
Publication
-
- 21 [biblio.ugent.be](#) <1 %
Internet Source
-
- 22 [doc-pak.undip.ac.id](#) <1 %
Internet Source
-
- 23 [hau.repository.guildhe.ac.uk](#) <1 %
Internet Source
-
- [repositorio-aberto.up.pt](#)

- Internet Source **24** <1 %
-
- Internet Source **25** rik.mrzp.rs <1 %
-
- Internet Source **26** www.researcherslinks.com <1 %
-
- Publication **27** Mahmoud A.O. Dawood, Marwa F. Abd El-Kader, Mona A. Farid, Mohamed F. Abd-Elghany, Mohamed Alkafafy, Hien Van Doan. "enhanced the growth, immune and antioxidative responses of European seabass ()", Annals of Animal Science, 2021 <1 %
-
- Publication **28** Mohammod Kamruzzaman Hossain, Sairatul Dahlianis Ishak, Shumpei Iehata, Mat Noordin Noordiyana et al. "Growth performance, fatty acid profile, gut, and muscle histomorphology of Malaysian mahseer, *Tor tambroides* post larvae fed short-term host associated probiotics", Aquaculture and Fisheries, 2022 <1 %
-
- 29** Soleimani, N.. "Dietary supplementation of fructooligosaccharide (FOS) improves the innate immune response, stress resistance, digestive enzyme activities and growth <1 %

performance of Caspian roach (*Rutilus rutilus*) fry", Fish and Shellfish Immunology, 201202

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches < 5 words