

DAFTAR PUSTAKA

1. Baradero M, Daydrit MW, Siswadi. Asuhan keperawatan klien gangguan ginjal. Jakarta: ECG; 2009.
2. Price SA WL. Faktor risiko gagal ginjal pada diabetes melitus. 2006;1(1).
3. Kemenkes RI. Hasil utama riset kesehatan dasar (Riskesdas) 2018.
4. Registry IR. 9 th Report of indonesian renal registry 2016. 2016;
5. PERNEFRI. 13th Annual report of indonesian renal registry 2020. Indonesia Renal Registry
6. Muttaqin A SK. Asuhan keperawatan gangguan sistem perkemihan. Jakarta: Salemba Medika; 2011.
7. Samuel Raj V V, Mangalvedhe P V, Shetty MS, Balakrishnan DC. Impact of exercise on fatigue in patients undergoing dialysis in a tertiary care hospital. Cureus. 2023;15(2).
8. Tim Pokja SDKI DPP PPNI. Standar diagnosis keperawatan Indonesia. Persat Perawat Indones. 2016;
9. Ju A, Strippoli GFM, Craig JC, Tong A, Saglimbene VM, Unruh ML. Interventions for fatigue in people with chronic kidney disease requiring dialysis. Cochrane Database Syst Rev. 2018;2018(8).
10. Parker Gregg L, Bossola M, Ostrosky Frid M, Susan HS. Fatigue in ckd epidemiology, pathophysiology, and treatment. Clin J Am Soc Nephrol.

2021;16(9):1445–55.

11. Wahida AZ, Rumahorbo H, Murtiningsih. The effectiveness of intradialytic exercise in ameliorating fatigue symptoms in patients with chronic kidney failure undergoing hemodialysis: A systematic literature review and meta-analysis. *J Taibah Univ Med Sci.* 2023;18(3):512–25.
12. Anggela S, Agustini N. Penerapan relaksasi otot progresif terhadap fatigue pada anak penyakit ginjal yang menjalani hemodialisis. *JONAH (Journal Nurs Homecare.* 2023;2(1):1–57.
13. Wallin H, Jansson E, Wallquist C, Hylander RB, Jacobson SH, Rickenlund A, et al. Aerobic exercise capacity is maintained over a 5-year period in mild-to-moderate chronic kidney disease: a longitudinal study. *BMC Nephrol.* 2020;21(1):1–11.
14. Prastiwi F, Wihastuti TA, Ismail DDSL. Factors associated with fatigue in chronic kidney disease patients undergoing hemodialysis: a systematic review. *J Kedokt Brawijaya.* 2021;31(4):260–8.
15. Firdaus ANT, Waluyo A, Jumaiyah W. Stretching exercise between the time of dialysis for fatigue scores of chronic kidney failure patient at the general hospital of ciamis district. *J Wacana Kesehat Univ Muhammadiyah Cirebon.* 2022;7(1):32–40.
16. Mohamed SHM. Effect of intradialytic exercise on fatigue, electrolytes level and blood pressure in hemodialysis patients: A randomized controlled trial. *J Nurs Educ Pract.* 2015;5(11):16–28.

17. Kurniawati BD, Wahyuni TD, Wicaksana AL. The effect of intradialytic exercise on fatigue-related chronic kidney disease: a case study of patient undergoing hemodialysis. *Open Access Maced J Med Sci.* 2022;10(C):290–3.
18. Mufidah Nurdesia, Mohammad Ali ZS. *Indonesian Journal of Physiotherapy.* Pengaruh aerobic exercise terhadap tekanan darah pada penderita hipertensi *Stud Lit.* 2021;1(1):1–5.
19. Kisner, Carolyn, Colby, Lynn Allen, Borstad J. *Therapeutic exercise : foundations and techniques.* Seventh Ed. A. Davis Company; 2018.
20. Daimo, M., Mandal, S., Mahmud, M., & Mathivanan D. Effect of aerobic exercise on blood pressure in men with hypertension: A randomized controlled study. *Turkish J Kinesiol.* 2020;6(1):32-39.
21. Rashid-Farokhi F, Pourdowlat G, Nikoonia MR, Behzadnia N, Kahkouee S, Nassiri AA, et al. Uremic pleuritis in chronic hemodialysis patients. *Hemodial Int.* 2013;17(1):94–100.
22. Wilkinson TJ, Watson EL, Gould DW, Xenophontos S, Clarke AL, Vogt BP, et al. Twelve weeks of supervised exercise improves self-reported symptom burden and fatigue in chronic kidney disease: A secondary analysis of the “ExTra CKD” trial. *Clin Kidney J.* 2019;12(1):113–21.
23. Sari Anggela, Allenidekania NA. Penerapan relaksasi otot progresif terhadap fatigue. 2023;2(1).
24. Muliani R, Muslim AR, Abidin I. Intradialytic exercise : flexibility terhadap skor fatigue pada pasien penyakit ginjal kronis yang menjalani hemodialisis

- intradialytic exercise : flexibility on fatigue scores in chronic kidney disease clients undergoing hemodialysis. *J Med Heal*. 2021;3(2):137–46.
25. Makiyah N, Sakitri G KA. The effect of intradialytic exercise on fatigue hemodialysis patients at RSUP dr. Soeradji Tirtonegoro Klaten. *Media Publ Penelit Prof Islam [Internet]*. 2017;15(1):58–64. Available from: <https://doi.org/10.26576/profesi.242>
 26. Ganik S. Pengaruh intradialytic exercise terhadap fatigue, Kadar hemoglobin dan tekanan darah pasien hemodialysis di RSUP Dr. Soradji Titonegoro Klaten. *Media Publ Penelit Prof Islam [Internet]*. 2018;1–17. Available from: <https://doi.org/%0A10.26576/profesi.242>
 27. Amilia y, Bakar A NI. The effect of intradialytic exercise with the peaceful end of life approach to fatigue of end stage renal disease patients who have dialysis. *Proc Surabaya Int Heal conf*. 2019;150–7.
 28. Djamaludin D, Chrisanto EY WM. Pengaruh latihan fisik terhadap penurunan fatigue pada pasien gagal ginjal kronik yang menjalani hemodialisa di RSUD dr. H. Abdul Moeloek Provinsi Lampung. *Malahayati Nurse J [Internet]*. 2020;667–76. Available from: <https://doi.org/10.33024/manuju.v2i4.1623>
 29. Lerdal, A., & Kottorp A. Psychometric properties of the fatigue severity Scale—Rasch analyses of individual responses in a norwegian stroke cohort. *International Journal of Nursing Studies*. 2011;48(10):1258–65.
 30. Pisoni RL, Young EW, Dykstra DM, Greenwood RN, Hecking E, Gillespie B, et al. Vascular access use in Europe and the United States: results from the

- DOPPS. *Kidney Int.* 2002;61(1):305–16.
31. Mohamed S. The effectiveness of an educational intervention on fatigue in hemodialysis patients: a randomized controlled trial. *IOSR J Nurs Heal Sci.* 2020;3(4):40–50.
 32. Grigoriou SS, Krase AA, Karatzaferi C, Giannaki CD, Lavdas E, Mitrou GI, et al. Long-term intradialytic hybrid exercise training on fatigue symptoms in patients receiving hemodialysis therapy. *Int Urol Nephrol* [Internet]. 2021;53(4):771–84. Available from: <https://doi.org/10.1007/s11255-020-02711-8>
 33. Amri K, Pangambam S, Bhutia KD. Effectiveness of intradialytic leg ergometry and stretching exercises on fatigue and muscle cramps among patients subjected to hemodialysis at tertiary care hospitals of sikkim. *Int J Heal Sci Res.* 2023;13(2):17–20.
 34. Kusuma H, Ropyanto CB, Hastuti YD, Hidayati W, Sujianto U, Setiawan D, et al. Modul pendampingan perawatan kesehatan mandiri dalam manajemen penyakit ginjal kronik-hipertensi. 2020; 2020.
 35. Suwitra K. Penyakit ginjal kronik. v. buku ajar ilmu penyakit dalam; 2009. 1035 p.
 36. Walklin CG, Young HML, Asghari E, Bhandari S, Billany RE, Bishop N, et al. The effect of a novel, digital physical activity and emotional well-being intervention on health-related quality of life in people with chronic kidney disease: trial design and baseline data from a multicentre prospective, wait-list

- randomised controlled . *BMC Nephrol.* 2023;24(1):1–17.
37. Khanifah AN. Gambaran pola tidur pada penderita penyakit ginjal kronik di ruang hemodialisa RS lavalette Kota Malang. 2022;
 38. Ratna, M. E., Windartik, E., & Pratiwi RM. Hubungan kadar hemoglobin (hb) dengan kelelahan (fatigue) pada pasien gagal ginjal kronis yang menjalani hemodialisa di ruang HD RSI Siti Hajar Sidoarjo. 2023;
 39. Joshwa, B., & Campbell ML. Fatigue in patients with chronic kidney disease: evidence and measures. *Nephrol Nurs J.* 2017;44(4).
 40. Artom M, Moss-Morris R, Caskey F, Chilcot J. Fatigue in advanced kidney disease. *Kidney Int [Internet].* 2014;86(3):497–505. Available from: <http://dx.doi.org/10.1038/ki.2014.86>
 41. Jacobson J, Ju A, Baumgart A, Unruh M, O’Donoghue D, Obrador G, et al. Patient perspectives on the meaning and impact of fatigue in hemodialysis: a systematic review and thematic analysis of qualitative studies. *Am J Kidney Dis.* 2019;74(2):179–92.
 42. Levine M. Introduction to clinical nursing(2nd ed.). In McEwen, M. and Wills, E. (Ed.). *Theoretical basis for nursing.* Lippincott Williams & Wilkins. 1973;
 43. Maryline E Parker. *Nursing theories and nursing practice.* Elsevier. 2005;4th Editio.
 44. Levine M. The conservation principles of nursing: Twenty years later. In George, J. (Ed.). *Nursing theories: the base for professional nursing practice.* Norwalk, Connect Applet Lange. 1989;

45. LC M. A Theory of health promotion for preterm infants based on levine's conservation model of nursing. *Nurs Sci Q.* 2004;17(3):260–6.
46. Kazancioğlu R. Risk factors for chronic kidney disease: An update. *Kidney Int Suppl.* 2013;3(4):368–71.
47. Novi TF A, Waluyo A, Jumaiyah W. Stretching exercise between the time of dialysis for fatigue scores of chronic kidney failure patient at the general hospital of ciamis district. 2022;7(1):32–40.
48. C J Caspersen, K E Powell and GMC. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Notes Queries.* 1985;s9-IX(228):365–6.
49. Tabibi MA, Cheema B, Salimian N, Corrêa H de L, Ahmadi S. The effect of intradialytic exercise on dialysis patient survival: a randomized controlled trial. *BMC Nephrol [Internet].* 2023;24(1):1–9. Available from: <https://doi.org/10.1186/s12882-023-03158-6>
50. Mailani F, Herien Y, Muthia R, Tumanggor RD, Huriani E. The experiences of patients with chronic kidney disease undergoing dialysis in managing the symptoms. *Malaysian J Med Heal Sci.* 2022;18(4):42–50.
51. Ma'ruf MR, Anggeria E, Siregar RF, Sinaga ES, Aminah, Rahmi IM, et al. Application of swedish massage therapy for lower extremities to alleviate muscle cramps in chronic kidney disease patients. *Futur Med.* 2024;3(4):4–10.
52. Leal-Martín J, Muñoz-Muñoz M, Keadle SK, Amaro-Gahete F, Alegre LM, Mañas A, et al. Resting oxygen uptake value of 1 metabolic equivalent of task

- in older adults: a systematic review and descriptive analysis. *Sport Med*
53. Adeel M, Lai CH, Wu CW, Kang JH, Liou JC, Chen HC, et al. Modeling of metabolic equivalents (Mets) during moderate resistance training exercises. *Appl Sci*. 2021;11(18):1–12.
 54. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR, Tudor-Locke C, et al. 2011 compendium of physical activities: A second update of codes and MET values. *Med Sci Sports Exerc*. 2011;43(8):1575–81.
 55. Martha Raile Alligood. *Nursing theories and their work*. Elsevier. 2014;8.
 56. Sun G, Hao R, Zhang L, Shi X, Hei K, Dong L, et al. The effect of hemodialysis on ocular changes in patients with the end-stage renal disease. *Ren Fail*. 2019;41(1):629–35.
 57. Johansen KL, Shubert T, Doyle J, Soher B, Sakkas GK, Kent-Braun JA. Muscle atrophy in patients receiving hemodialysis: effects on muscle strength, muscle quality, and physical function. *Kidney Int*. 2003;63(1):291–7.
 58. Nursalam. *Konsep dan penerapan metodologi penelitian ilmu keperawatan*. Jakarta: salemba medika; 2008. 89–93 p.
 59. Amini E, Goudarzi I, Masoudi R, Ahmadi A, Momeni A. Effect of progressive muscle relaxation and aerobic exercise on anxiety, sleep quality, and fatigue in patients with chronic renal failure undergoing hemodialysis. *Int J Pharm Clin Res*. 2016;8(12):1634–9.
 60. Acaster S, Dickerhoof R, DeBusk K, Bernard K, Strauss W, Allen LF. Qualitative and quantitative validation of the FACIT-fatigue scale in iron

- deficiency anemia. *Health Qual Life Outcomes*. 2015;13(1).
61. Sihombing JP, Hakim L, Andayani TM, Irijanto F. Validation of Indonesian version of Facit fatigue scale questionnaire in chronic kidney disease (CKD) patients with routine hemodialysis. *Indones J Clin Pharm [Internet]*. 2016 Dec 1;5(4):231–7. Available from: <http://jurnal.unpad.ac.id/ijcp/article/view/13486>
 62. Kuningas K, Inston N. Age is just a number: Is frailty being ignored in vascular access planning for dialysis? *J Vasc Access*. 2022;23(2):192–7.
 63. Llisterri JL, Micó-Pérez RM, Velilla-Zancada S, Rodríguez-Roca GC, Prieto-Díaz MÁ, Martín-Sánchez V, et al. Prevalence of chronic kidney disease and associated factors in the Spanish population attended in primary care: Results of the IBERICAN study. *Med Clínica (English Ed)*. 2021;156(4):157–65.
 64. Carrero JJ, Hecking M, Chesnaye NC, Jager KJ. Sex and gender disparities in the epidemiology and outcomes of chronic kidney disease. *Nat Rev Nephrol [Internet]*. 2018;14(3):151–64. Available from: <http://dx.doi.org/10.1038/nrneph.2017.181>
 65. Motiei M, Attarchi M, Ramezanzadeh E. The effect of workability-related factors in patients with end-stage kidney disease undergoing hemodialysis. *BMC Nephrol [Internet]*. 2024;25(1). Available from: <https://doi.org/10.1186/s12882-024-03904-4>
 66. de Jong RW, Boezeman EJ, Chesnaye NC, Bemelman FJ, Massy ZA, Jager KJ, et al. Work status and work ability of patients receiving kidney replacement therapy: results from a European survey. *Nephrol Dial Transplant*.

2022;37(10):2022–33.

67. Sajjadi SL, Ghafourifard M, Khosroshahi HT. The effect of individualized education on learning needs of patients undergoing hemodialysis: a randomized controlled clinical trial. *BMC Nephrol.* 2024;25(1).
68. Rini IS, Rahmayani T, Sari EK, Lestari R. Differences in the quality of life of chronic kidney disease patients undergoing hemodialysis and continuous ambulatory peritoneal dialysis. *J Public health Res.* 2021;10(2):301–5.
69. Devi S, Rahman S. The relationship of duration undergoing hemodialysis therapy and the quality of life of patients with chronic kidney disease at the rasyida kidney hospital. *ARC J Public Heal Community Med.* 2020;5(2):15–21.
70. Duan, J. Y., G. C. Duan, C. J. Wang, D. W. Liu, Y. J. Qiao, S. K. Pan, et al. Prevalence and risk factors of chronic kidney disease and diabetic kidney disease in a central chinese urban population: a cross-sectional survey. *BMC Nephrol.* 2020;21(115):1–13.
71. Balamuthusamy S, Miller LE, Clynes D, Kahle E, Knight RA, Conway PT. American association of kidney patients survey of patient preferences for hemodialysis vascular access. *J Vasc Access.* 2020;21(2):230–6.
72. Di D, Kitamura K, Ridwan M, Putra C. Naskah publikasi. 2017.
73. Sebayang ANO, Hidayat NA. Arteriovenous shunt as the best hemodialysis access in chronic kidney disease (CKD) patients: a literature review. *J Indones Vasc Access.* 2021;1(1):1–3.

74. De Clerck D, Bonkain F, Cools W, Van Der Niepen P. Vascular access type and mortality in haemodialysis: A retrospective cohort study. *BMC Nephrol.* 2020;21(1):1–7.
75. Hasibuan SF. Profile of anemia and the relationship between hemoglobin levels and quality of life in end-stage chronic kidney disease patients undergoing chronic hemodialysis at hasan sadikin hospital 2021-2022. *Indones J Kidney Hypertens.* 2024;1(1):17–23.
76. Yin P, Wu Q, Shou L, Dong X. Risk factors for anemia in patients with chronic kidney disease: A protocol for systematic review and meta-analysis. *Med (United States).* 2021;100(40):E27371.
77. Malini H, Forwaty E, Cleary M, Visentin D, Oktarina E, Lenggogeni DP. The effect of intradialytic range of motion exercise on dialysis adequacy and fatigue in hemodialysis patients. *J Nurs Res.* 2022;30(4):E221.
78. Lakshmi Y, D S, Varughese S. The Effect of intradialytic aerobic exercise on dialysis parameters and fatigue in hemodialysis patients: a non-randomized interventional study. *Cureus.* 2024;16(6).
79. Knoop V, Cloots B, Costenoble A, Debain A, Vella Azzopardi R, Vermeiren S, et al. Fatigue and the prediction of negative health outcomes: A systematic review with meta-analysis. *Ageing Res Rev.* 2021;67(September 2020).
80. Sridharan S, Vilar E, Ramanarayanan S, Davenport A, Farrington K. Energy expenditure estimates in chronic kidney disease using a novel physical activity questionnaire. *Nephrol Dial Transplant.* 2022;37(3):515–21.

81. Odetunde MO, Makinde AF, Jimoh OM, Mbada CE, Odumosu FN, Fatoye F. Physical activity , fatigue severity , and health - related quality of life of community - dwelling stroke survivors : a cross - sectional study. Bull Fac Phys Ther
82. Bossola M, Di Stasio E, Antocicco M, Panico L, Pepe G, Tazza L. Fatigue is associated with increased risk of mortality in patients on chronic hemodialysis. Nephron. 2015;130(2):113–8.
83. Vestergaard S, Nayfield SG, Patel K V., Eldadah B, Cesari M, Ferrucci L, et al. Fatigue in a representative population of older persons and its association with functional impairment, functional limitation, and disability. Journals Gerontol - Ser A Biol Sci Med Sci. 2009;64(1):76–82.
84. Koyama H, Fukuda S, Shoji T, Inaba M, Tsujimoto Y, Tabata T, et al. Fatigue is a predictor for cardiovascular outcomes in patients undergoing hemodialysis. Clin J Am Soc Nephrol. 2010;5(4):659–66.
85. Rhee SY, Song JK, Hong SC, Choi JW, Jeon HJ, Shin DH, et al. Intradialytic exercise improves physical function and reduces intradialytic hypotension and depression in hemodialysis patients. Korean J Intern Med. 2019;34(3):588–98.
86. Florent Besnier, Eric Laruelle, Sandrine Genestier, Sophie Gie' a B, , Ce' cile Vigneau b C, Carre F. Effects of exercise training on ergocycle during hemodialysis in patients with end stage renal disease: Relevance of the anaerobic threshold intensity. Nephrol Ther. 2012;231–7.
87. Handling C, Cheng AJ, Place N. Molecular basis for exercise-induced fatigue:

The importance of strictly controlled cellular Ca²⁺ handling. *Cold Spring Harb Perspect Med.* 2018;8:a029710.

88. Zhao M, Xiao M, Tan Q, Lyu J, Lu F. The effect of aerobic exercise on oxidative stress in patients with chronic kidney disease: a systematic review and meta-analysis with trial sequential analysis. *Ren Fail* [Internet]. 2023;45(2). Available from: <https://doi.org/10.1080/0886022X.2023.2252093>
89. Zhao R, Wu R, Jin J, Ning K, Wang Z, Yi X, et al. Signaling pathways regulated by natural active ingredients in the fight against exercise fatigue-a review. *Front Pharmacol.* 2023;14(December):1–14.
90. Nakoui N, Ilbeigi S, Ahmadi MM, Saber A. Comparison of the effect of aerobic and resistance training on fatigue, quality of life and biochemical factors in hemodialysis patients. *Sci Rep.* 2025;15(1):1–11.
91. Hatef M, Esmacili R, Mousavinasab N, Madani Z, Spahbodi F, Shafipour V. Effect of intradialytic and home-based walking exercises on physical function and dialysis adequacy in hemodialysis patients. *J Maz Univ Med Sci.* 2017;27(154):83–93.
92. Chang Y, Cheng SY, Lin M, Gau FY, Chao YFC. The effectiveness of intradialytic leg ergometry exercise for improving sedentary life style and fatigue among patients with chronic kidney disease: A randomized clinical trial. *Int J Nurs Stud* [Internet]. 2010;47(11):1383–8. Available from: <http://dx.doi.org/10.1016/j.ijnurstu.2010.05.002>
93. Ritianingsih N, Nawati N, Nurhayati F. The energy conservation strategies can

improve self care management of chronic kidney disease patients with hemodialysis. *Indones J Glob Heal Res.* 2023;5(2):319–26.

94. Pedersen H, Fimland MS, Schoenfeld BJ, Iversen VM, Cumming KT, Jensen S, et al. A randomized trial on the efficacy of split-body versus full-body resistance training in non-resistance trained women. *BMC Sports Sci Med Rehabil [Internet]*. 2022;14(1):1–9. Available from: <https://doi.org/10.1186/s13102-022-00481-7>
95. Mohseni R, Zeydi AE, Ilali E, Hajbaghery MA and AM. The effect of intradialytic aerobic exercise on dialysis efficacy in hemodialysis patients: A randomized controlled trial. *Oman Med J.* 2013;28(5):345–9.