

DAFTAR PUSTAKA

- Adnan, M., Bibi, R., Azizullah, A., Andaleeb, R., Mussarat, S., Tariq, A., ... & Begum, S. (2015). Ethnomedicinal plants used against common digestive problems. *African Journal of Traditional, Complementary and Alternative Medicines*, 12(5), 99-117.
- Ahmed, M., Singh, M. N., Bera, A. K., Bandyopadhyay, S., & Bhattacharya, D. (2011). Molecular basis for identification of species/isolates of gastrointestinal nematode parasites. *Asian Pacific Journal of Tropical Medicine*, 4(8), 589-593.
- Uttra, A. M., Ahsan, H., Hasan, U. H., & Chaudhary, M. A. (2018). Traditional medicines of plant origin used for the treatment of inflammatory disorders in Pakistan: A review. *Journal of Traditional Chinese Medicine*, 38(4), 636-656.
- Argus, G. W. (1995). Salicaceae Willow Family: Part Two: *Salix* L. Willow. *Journal of the Arizona-Nevada Academy of Science*, 29(1), 39–62.
- Argus, G. W. (1997). Infrageneric Classification of *Salix* (Salicaceae) in the New World. *Systematic Botany Monographs*, 52, 1–121.
- Atawodi, S.E., Atawodi, J.C. and Dzikwi, A.A. 2010. Polymerase Chain Reaction: Theory, Practice and Application. *Sahel Medical Journal*, vol. 13, no. 2.
- Aulia, S. L., Suwignyo, R. A., & Hasmeda, M. (2021). Optimasi Suhu Annealing untuk Amplifikasi Dna Padi Hasil Persilangan Varietas Tahan Terendam dengan Metode Polymerase Chain Reaction. *Sainmatika: Jurnal Ilmiah Matematika dan Ilmu Pengetahuan Alam*, 18(1), 44-54.
- Baldovini, N., Delasalle, C., & Joulain, D. (2011). Phytochemistry of the heartwood from fragrant Santalum species: a review. *Flavour and Fragrance Journal*, 26(1), 7-26.
- Batiha, G. E. S., Beshbishy, A. M., Guswanto, A., Nugraha, A., Munkhjargal, T., M. Abdel-Daim, M., ... & Igarashi, I. (2020). Phytochemical characterization and chemotherapeutic potential of *Cinnamomum verum* extracts on the multiplication of protozoan parasites in vitro and in vivo. *Molecules*, 25(4), 996.
- Belyaeva, I. V., Epantchintseva, O. V., Govaerts, R. H., McGinn, K., Hunnex, J., & Kuzovkina, Y. A. (2018). The application of scientific names to plants

in cultivation: *Salix vitellina* L. and related taxa (Salicaceae). *Skvortsovia*, 4(2), 42-70.

Betz, O. (2006). Ecomorphology: Integration of form, function, and ecology in the analysis of morphological structures. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie*, 15, 409-416.

Calonje, M., Martín-Bravo, S., Dobeš, C., Gong, W., Jordon-Thaden, I., Kiefer, C., Kiefer, M., Paule, J., Schmickl, R. and Koch, M.A. (2009). Non-coding nuclear DNA markers in phylogenetic reconstruction. *Plant Systematics and Evolution*, 282, 257-280. markers in phylogenetic reconstruction. *Plant Syst Evol* 282, 257–280

Cervantes, E. and de Diego, J. G. 2010. Morphological description of plants. New perspectives in development and evolution. *Int. J. Pl. Dev. Biol.* Vol. 4, No. 1, pp. 68-71.

Collard, B. C., Jahufer, M. Z. Z., Brouwer, J. B., & Pang, E. C. K. (2005). An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: the basic concepts. *Euphytica*, 142, 169-196.

Dimitriou, I., & Aronsson, P. (2010). Landfill leachate treatment with willows and poplars—Efficiency and plant response. *Waste management*, 30(11), 2137-2145.

Doyle, J. J. (1990). Isolation of plant DNA from fresh tissue. *Focus*, 12, 13-15.

eFloras. (2008). *Flora of China*, Vol. 23, *Arecaceae*, *Arenga*. pp. 151-152. Available from: http://www.efloras.org/florataxon.aspx?flora_id=2&taxon_id=250095411 (accessed: 25 Juni 2020).

El-Shemy, H. A., Aboul-Enein, A. M., Aboul-Enein, K. M., & Fujita, K. (2007). Willow leaves' extracts contain anti-tumor agents effective against three cell types. *Plos one*, 2(1), e178.

Elya, B. E. R. N. A., Puspitasari, N. U. R. A. I. N. I., & Sudarmin, A. C. (2017). Antioxidant activity and inhibition of lipoxygenase activity ethanol extract of endosperm *Arenga pinnata* (Wurmb) Merr. *Asian Journal of Pharmaceutical and Clinical Research*, 10 (Special Issue October), 76-80.

- Fabbri, A., Hormaza, J. I., & Polito, V. S. (1995). Random amplified polymorphic DNA analysis of olive (*Olea europaea* L.) cultivars. *Journal of the American Society for Horticultural Science*, *120*(3), 538-542.
- Farias, A. P. P., Monteiro, O. D. S., da Silva, J. K. R., Figueiredo, P. L. B., Rodrigues, A. A. C., Monteiro, I. N., & Maia, J. G. S. (2020). Chemical composition and biological activities of two chemotype-oils from *Cinnamomum verum* J. Presl growing in North Brazil. *Journal of Food Science and Technology*, *57*, 3176-3183.
- Fortini, P., Di Marzio, P., Guarrera, P. M., & Iorizzi, M. (2016). Ethnobotanical study on the medicinal plants in the Mainarde Mountains (central-southern Apennine, Italy). *Journal of Ethnopharmacology*, *184*, 208-218.
- Fristseh, P., & Rieseberg, L. H. (1996). The use of random amplified polymorphic DNA (RAPD) in conservation genetics. Molecular genetic approaches in conservation. Univ. Press.
- Gariyban, L., & Avashia, N. (2013). Research techniques made simple: polymerase chain reaction (PCR). *The Journal of investigative dermatology*, *133*(3), e6.
- Gulyuz, H., Fazilat, S., & Komila, S. (2023). Information About *Salix Matsudana*. *World Journal of Agriculture and Urbanization*, *2*(11), 6–8. <https://doi.org/10.51699/wjau.v2i11.11>
- Gunawardena, D., Karunaweera, N., Lee, S., van Der Kooy, F., Harman, D. G., Raju, R., ... & Münch, G. (2015). Anti-inflammatory activity of cinnamon (*C. zeylanicum* and *C. cassia*) extracts—identification of E-cinnamaldehyde and o-methoxy cinnamaldehyde as the most potent bioactive compounds. *Food & function*, *6*(3), 910-919.
- Halimatussakdiah, H., Amna, U., & Wahyuningsih, P. (2018). Preliminary phytochemical analysis and larvicidal activity of edible fern (*Diplazium esculentum* (retz.) sw.) extract against culex. *Jurnal Natural*, *18*(3), 141-147.
- Harbourne, N., Marete, E., Jacquier, J. C., & O'Riordan, D. (2009). Effect of drying methods on the phenolic constituents of meadowsweet (*Filipendula ulmaria*) and willow (*Salix alba*). *LWT-Food Science and Technology*, *42*(9), 1468-1473.

- Hikmatyar, M. F., & Royani, J. I. (2015). Isolasi dan Amplifikasi DNA Keladi Tikus (*Thyponium flagelliform*) untuk Identifikasi Keragaman Genetik. *Jurnal Bioteknologi dan Biosains Indonesia*, 2(2), 42-48.
- Huang, J., Liu, Y., Zhu, T., & Yang, Z. (2021). The asymptotic behavior of bootstrap support values in molecular phylogenetics. *Systematic biology*, 70(4), 774-785. doi: 10.1093/sysbio/syaa100. PMID: 33377913.
- Isebrands, J. G., and Richardson, J. (2014). *Poplars and willows: trees for society and the environment*. Boston, USA: The Food and Agriculture Organization of the United Nations and CABI, 634.
- Ismail, Z., Ismail, N., Jaafar, L. 1999. *Malaysian Herbal Monograph*, Vol. 1. Kuala Lumpur: Malaysian Monograph Committee.
- IUCN. 2007. IUCN Red List of Threatened Species. <https://www.iucnredlist.org/> diakses pada tanggal 28 November 2021 pada pukul 1.46.
- Jeanson, M. L., Labat, J. N., & Little, D. P. (2011). DNA barcoding: a new tool for palm taxonomists?. *Annals of botany*, 108(8), 1445-1451. doi: 10.1093/aob/mcr158.
- Jiang, G. L. (2013). Molecular markers and marker-assisted breeding in plants. *Plant breeding from laboratories to fields*, 3, 45-83.
- Jorgensen, R., Snyder, C., & Jones, J. D. (1987). T-DNA is organized predominantly in inverted repeat structures in plants transformed with *Agrobacterium tumefaciens* C58 derivatives. *Molecular and General Genetics MGG*, 207(2), 471-477.
- Liang, Z., Pan, C., Zhang, Z., Wang, Y., Liu, M., Chen, P., & Chen, J. (2023). Effects of electrophoresis buffer, electrophoresis time, and annealing temperature on the quality of SSR molecular detection in rice varieties. *Cogent Food & Agriculture*, 9(2), 2275414. doi: <https://doi.org/10.1080/23311932.2023.2275414>
- Keoleian, G. A., & Volk, T. A. (2005). Renewable energy from willow biomass crops: life cycle energy, environmental and economic performance. *BPTS*, 24(5-6), 385-406.
- Klasnja, B., Kopitovic, S., & Orlovic, S. (2002). Wood and bark of some poplar and willow clones as fuelwood. *Biomass and Bioenergy*, 23(6), 427-432.

- Kool, A., de Boer, H. J., Krüger, Å., Rydberg, A., Abbad, A., Björk, L., & Martin, G. (2012). Molecular identification of commercialized medicinal plants in Southern Morocco. *PloS one*, 7(6), e39459.
- Kristanti, A. N. (Ed.). (2019). *Fitokimia*. Airlangga University Press.
- Kumar, R., Anjum, N., & Tripathi, Y. C. (2015). Phytochemistry and pharmacology of *Santalum album* L.: a review. *World Journal of Pharmaceutical Research*, 4(10), 1842-1876.
- Kusmana, C., & Hikmat, A. (2015). Keanekaragaman hayati flora di Indonesia. *Jurnal Pengelolaan Sumberdaya Alam dan Lingkungan (Journal of Natural Resources and Environmental Management)*, 5(2), 187-187.
- Lauron-Moreau, A., Pitre, F. E., Argus, G. W., Labrecque, M., & Brouillet, L. (2015). Phylogenetic relationships of american willows (*Salix* L., Salicaceae). *PLoS One*, 10(4), e0121965.
- Li, X., Yang, Y., Henry, R. J., Rossetto, M., Wang, Y., & Chen, S. (2015). Plant DNA barcoding: from gene to genome. *Biological Reviews*, 90(1), 157-166.
- Mabberley, D. J. (2008). *Mabberley's Plant-book: a portable dictionary of plants, their classifications and uses*(No. Ed. 3). Cambridge university press.
- Mahdi, J. G. (2010). Medicinal potential of willow: A chemical perspective of aspirin discovery. *Journal of Saudi Chemical Society*, 14(3), 317-322.
- Maretha, D. E., Hapida, Y., & Nugroho, Y. A. T. (2020). Pemanfaatan Air Nira Tanaman Aren (*Arenga Pinnata* Merr) Menjadi Gula Semut. *Noer Fikri Offset. Palembang*.
- Miles, D. H. (1985). *A Guide to Biologically Active Plants Constituent*. Department of Chemistry, University of Florida, USA.
- Misra, B. B., & Dey, S. (2013). Developmental variations in sesquiterpenoid biosynthesis in East Indian sandalwood tree (*Santalum album* L.). *Trees*, 27, 1071-1086.
- Mondini, L., Noorani, A., & Pagnotta, M. A. (2009). Assessing plant genetic diversity by molecular tools. *Diversity*, 1(1), 19-35.
- Muhajidin, Sutrisno, Latifah, D., et al. (2003). *Aren Budi Daya dan Prospeknya*. Bogor : Pusat Konservasi Tumbuhan Kebun Raya Bogor.

- Musinguzi, C., Rutebemberwa, E., Namale, L., & Dahal, A. (2017). 203: The Relationship between Leadership Styles and Health Worker Motivation, Teamwork and Job Satisfaction in Health Facilities in Uganda. *BMJ Open*, 7(Suppl 1).
- Muthuswamy, S., Rupasinghe, H. P. V., & Stratton, G. W. (2008). Antimicrobial effect of cinnamon bark extract on *Escherichia coli* O157: H7, *Listeria innocua* and fresh-cut apple slices. *Journal of Food Safety*, 28(4), 534-549.
- Muzzalupo, I., Fodale, A., Caravita, M.A., et al. (2007). Genetic relationships of Sicilian olive cultivars using RAPD markers. *Adv. Hort. Sci.*, vol. 21, no. 1, pp. 35-40.
- Muzzazinah. (2017). Metode filogenetik pada Indigofera. *Prosiding Seminar Nasional Pendidikan Biologi dan Biologi*, <http://seminar.uny.ac.id/sembiouny2017/prosiding/metode-filogenetik-pada-indigofera>.
- Nei, M., & Kumar, S. (2000). *Molecular evolution and phylogenetics*. Oxford university press.
- Nessa, F., Ismail, Z., & Mohamed, N. (2004a). Isolation of sterols and hydrocarbons from the leaves of *Blumea balsamifera* DC. *ACGC CHEMICAL RESEARCH COMMUNICATIONS*, 9-15.
- Nessa, F., Ismail, Z., Mohamed, N., & Haris, M. R. H. M. (2004). Free radical-scavenging activity of organic extracts and of pure flavonoids of *Blumea balsamifera* DC leaves. *Food Chemistry*, 88(2), 243-252.
- Nguyen, M. T. T., Awale, S., Tezuka, Y., Le Tran, Q., Watanabe, H., & Kadota, S. (2004). Xanthine oxidase inhibitory activity of Vietnamese medicinal plants. *Biological and Pharmaceutical Bulletin*, 27(9), 1414-1421.
- Noletto-Dias, C., Ward, J. L., Bellisai, A., Lomax, C., & Beale, M. H. (2018). Salicin-7-sulfate: A new salicinoid from willow and implications for herbal medicine. *Fitoterapia*, 127, 166-172.
- Norikura, T., Kojima-Yuasa, A., Shimizu, M., Huang, X., Xu, S., Kametani, S., Rho, S.N., Kennedy, D.O. and Matsui-Yuasa, I. (2008). Anticancer activities and mechanisms of *Blumea balsamifera* extract in hepatocellular carcinoma cells. *The American Journal of Chinese Medicine*, 36(02), 411-424.

- Nurmayulis, N., Susiyanti, S., Isminingsih, S., & Sari, R. P. (2021). Identifikasi morfologi tanaman aren asal Kabupaten Lebak. *Jurnal Agroekoteknologi*, 13(2), 179-190.
- Orwa, C., Mutua, A., et al. (2009). *Agroforestry Database: a tree reference and selection guide* version 4.0. Kenya : World Agroforestry Centre.
- Poczai, P., & Hyvönen, J. (2010). Nuclear ribosomal spacer regions in plant phylogenetics: problems and prospects. *Molecular biology reports*, 37(4), 1897–1912.
- Pokorny, J. (1991). Natural antioxidants for food use. *Trends in Food Science and Technology*, vol. 9, pp. 223–227.
- Porter, C. H., & Collins, F. H. (1991). Species-diagnostic differences in a ribosomal DNA internal transcribed spacer from the sibling species *Anopheles freeborni* and *Anopheles hermsi* (Diptera: Culicidae). *The American journal of tropical medicine and hygiene*, 45(2), 271-279.
- Pramono, E. (2002). The commercial use of traditional knowledge and medicinal plants in Indonesia. *Multi-Stakeholder Dialogue on Trade, Intellectual Property and Biological Resources in Asia held at BRAC Centre for Development Management, Rajendrapur, Bangladesh from April 19-21*.
- Putri, I. D. A. E. W., I Gusti Ayu Dewi R., et al. (2021). Analisis fitokimia nira Aren dan tuak Aren (*Arenga pinnata* (Wurmb) Merr.). *Jurnal Medika Udayana*, Vol. 10, No. 6.
- Ragasa, C.Y., Kristin, C. Co, A. L., and Rideout, J. A. (2005). Antifungal metabolites from *Blumea balsamifera*. *Natural Product Research*, vol. 19, no. 3, pp. 231–237.
- Saller, R., Melzer, J., and Felder, M. (2008). Pain relief with a proprietary extract of Willow bark in rheumatology. *Swiss Journal of Integrative Medicine*. Vol. 20, No. 3, pp. 156-162.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). *Molecular Cloning : A laboratory Manual*. 2nd Edition. Cold Spring Harbor Laboratory Press. USA.
- Sari, A & Anwar, A & Dwipa, I & Hervani, D. (2021). Morphological Characteristics of Sugar Palm [*Arenga Pinnata* Merr.] Seedling Growth Based on Cotyledon Petiole Position. IOP Conference Series: Earth and Environmental Science. 741.

- Saxena, M., Saxena, J., Nema, R., Singh, D., and Gupta, A. (2013). Phytochemistry of Medicinal Plants. *Journal of Pharmacognosy and Phytochemistry*, vol. 1, no. 6
- Shimin, C. (2004). Application of PCR in detection of food microorganism. *Journal of Handan agricultural college*, vol. 21, no. 4, pp. 23-25.
- Sourav, M. S. H., & Anwar, K. B. (2020). Arenga westerhoutii Griff. and Livistona jenkinsiana Griff., two new species of wild palms for Bangladesh.
- Stuessy, T. F. (2009). *Plant Taxonomy: the Systematic evaluation of comparative data*. New York : Columbia University Press.
- Sudan, R., Bhagat, M., Gupta, S., & Devi, T. (2013). Comparative analysis of cytotoxic and antioxidant potential of edible Cinnamomum verum (bark) and Cinnamomum tamala (Indian bay leaf). *Free Radicals and Antioxidants*, pp.1-4.
- Suffredini, I. B., Sader, H. S., Gonçalves, A. G., Reis, A. O., Gales, A. C., Varella, A. D., & Younes, R. N. (2004). Screening of antibacterial extracts from plants native to the Brazilian Amazon Rain Forest and Atlantic Forest. *Brazilian journal of medical and biological research*, 37, 379-384.
- Swasti, E. (2007). *Introduction to Plant Breeding*. Padang: Department of Agriculture, Faculty of Agriculture, University of Andalas. pp. 144.
- Tawfeek, N., Mahmoud, M. F., Hamdan, D. I., Sobeh, M., Farrag, N., Wink, M., & El-Shazly, A. M. (2021). Phytochemistry, pharmacology and medicinal uses of plants of the genus Salix: An updated review. *Frontiers in pharmacology*, 12, 593856.
- Tjitrosoepomo, G. (2007.) *Morfologi Tumbuhan*. Yogyakarta : Gadjah Mada University Press.
- Vasundhara, M., Thara, B. S., Radhika, B., Ashwini Jayaram, A. J., & Priyanka, R. (2015). Assessment of Indian Sandal wood (Santalum album L.,) seeds for seed oil production and fatty acid methyl esters, Vol. 4, No. 11, pp. 1416-1425.
- White, & Bruns, Tom & Lee, Steven & Taylor, John. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In PCR protocols a guide to methods and applications, 315–322. Academic Press, San Diego.

- Yunos, N.J., Sulaiman, A., Yusuf, N.A., and Ghazali, D.R. (2018). Physicochemical Properties and Enhanced Antioxidant Activity of *Arenga Pinnata* Sap Through Different Processing Methods. *Solid State Science and Technology*, vol. 26, no. 1, pp. 253-262.
- Zuhud, E.A.M. (2009). The potential of tropical forest as the buffer for natural medicine material for the nation's health. *Jurnal Bahan Alam Indonesia*, vol. 6, no. 6, pp. 227-232.