

Jurnal 3 - Deep Breathing Relaxation Techniques Improve Emotional Control on Tuberculosis Patients

by Meidiana Dwidiyanti

Submission date: 03-Aug-2021 11:12AM (UTC+0700)

Submission ID: 1627217038

File name: echniques_Improve_Emotional_Control_on_Tuberculosis_Patients.pdf (484.22K)

Word count: 2271

Character count: 12327

7

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/324238424>

Deep Breathing Relaxation Techniques Improve Emotional Control on Tuberculosis Patients

Article in *International Journal of Public Health Science (IJPHS)* · September 2016

DOI: 10.11591/ijphs.v5i3.4803

CITATIONS

0

READS

684

3 authors, including:



Meidiana Dwidiyanti

Universitas Diponegoro

4

31 PUBLICATIONS 20 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



mindfulness [View project](#)

3

All content following this page was uploaded by Meidiana Dwidiyanti on 30 September 2018.

The user has requested enhancement of the downloaded file.

Deep Breathing Relaxation Techniques Improve Emotional Control on Tuberculosis Patients

Domianus Namuwali¹, Fery Agusman Motuho Mendrofa², Meidiana Dwidiyanti³

^{1,3} Master Student of Nursing at Diponegoro University, Indonesia

² Department of Nursing Sciences, Karya Husada Institute of Health Science, Indonesia

Article Info

Article history:

Received Jun 17, 2016

Revised Aug 13, 2016

Accepted Aug 20, 2016

Keyword:

Emotional Control

Patient

Relaxation

Tuberculosis

ABSTRACT

Tuberculosis is an infectious disease caused by Tuberculosis Mycobacterium. Based on WHO report in 2014, the world population suffering from tuberculosis were 9.6 million people while in Indonesia it was amounted 324 539 people. Tuberculosis patients have a tendency to experience emotional disturbance due to the illness. A deep breathing relaxation is a nursing action controlling emotions of tuberculosis patients. The study design is quasi-experimental design with one group pre-test-post-test. Data were collected by using a questionnaire adapted from Gross and John (2003). Data were analyzed using paired t test. The results of this research is deep breathing relaxation technique is effective to control emotions of tuberculosis patients with p value=0,001.

1

Copyright © 2016 Institute of Advanced Engineering and Science.
All rights reserved.

Corresponding Author:

Domianus Namuwali,
Master Student of Nursing,
Diponegoro University,

Jalan Adam Malik No 126, Waingapu, Sumba Timur, Nusa Tenggara Timur, Indonesia.

Email : domianus2012@gmail.com

20

1. INTRODUCTION

18
Based on the WHO report in 2014 it was estimated that there were 9.6 million people worldwide who suffered from tuberculosis. Indonesia ranks third as the most populous country suffer tuberculosis by the number of tuberculosis in 2014, that was 324 539[1]. Patients with tuberculosis have a tendency to experience emotional disturbances. Results of research conducted by Paulo B.X & Bruno P states that 44.4% of tuberculosis patients experiencing emotional distress [2].

Patients who are undergoing treatment are necessary to get emotional control in order not to aggravate the patient's condition. Emotional control is an attempt to hide and to calm the emotions of expressing negative emotions such as anger, depression and anxiety. Control emotions affect the adjustment of disease through adaptive processes by using strategies of coping with stress [3].

A nurse is one of the team that provides health care in patients with tuberculosis that has a central role to provide emotional support for patients during illness and treatment [4]. One of the treatments to reduce emotional disorders in patients with tuberculosis is deep breathing relaxation. Relaxation that is performed for 5 to 10 minutes can improve blood flow throughout the body so that the body becomes more relaxed, and in the end it can relieve pain and mental stress in patients [5].

2. RESEARCH METHOD

15
The method used in this research was quasi-experimental. The study design was one group for pretest and posttest without a control group. The population in this study was all patients with pulmonary tuberculosis in Public Health Center (BKPM). There were 31 samples in this study. Inclusion criteria in this

1

Journal homepage: <http://iaesjournal.com/online/index.php/IJPHS>

study as follows: tuberculosis patients that were registered in Magelang BKPM, Patients under early and advanced stage of treatment, aged 15-60 years old, patients who do not experience mental illness, able to read and write and willing to be investigated. Exclusion criteria were patients with shortness of breath, extra pulmonary tuberculosis, complications of HIV/AIDS and diabetes mellitus.

Sampling was conducted by purposive sampling, the sample that is based on the intention and purpose as determined by researcher. The data collection was conducted using questionnaires adopted from Gross and John, which consists of 10 questions. Results of validity and reliability tests towards this questionnaire showed that only seven questions valid for the research, therefore, this study only used valid and reliable questionnaire. Pre test data was collected in the first week and then the respondents were trained in breath relaxation techniques at the first, the second and third weeks. Post test was conducted at the third week. Data analysis was performed using paired t test if the data normally distributed, but if it is not then it will be tested with Wilcoxon test [6].

3. RESULTS AND ANALYSIS

Characteristics of respondents presented in Table 1. The characteristics included gender, age, education, and occupation. In Table 1 it is known that most male respondents 18 (58.1%); the largest age group is the 26-45 years old, 15 (48.4%); most education of respondents is High School 17 (54.8%); and most respondents were private employees that was 9 (29%).

Table 1. Characteristics of Respondents

Characteristics of respondents	Frequency	Percent
Gender		
a. Male	18	58.1
b. Female	13	41.9
Age		
a. 15-25 years old	8	25.8
b. 26-45 years old	15	48.4
c. 47-60 years old	8	25.8
Education		
a. Elementary School	5	16.1
b. Junior High School	9	29.0
c. High school	17	54.8
Occupation		
a. Private employees	9	29.0
b. Entrepreneur	5	16.1
c. Labor	4	12.9
d. Housewife	3	9.7
e. Student	4	12.9
f. Etc.	2	6.5
g. Does not work	4	12.9

Sources: Primary data, in 2016.

3.1. Normality data test

In Table 2, the result of Shapiro-Wilk normality test shows significance values for the pre-test and posttest is > 0.05 . Thus, it can be concluded that the group of pre-test and post test data was normally distributed so that the analysis of the data used the paired t test.

Table 2. Test of Data Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	Df	Sig.
pretest	.175	31	.017	.941	31	.087
posttest	.075	31	.200 ^a	.979	31	.787

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

3.2. Emotion control

Table 3 presents the result of bivariate test. In Table 3 it is known that there are significantly differences in the average (mean) before and after relaxation, with the value of CI 95% between 5.65 to 10.28.

12

Table 3. Comparison of Mean and Standard Deviation Before and After the Deep Breath Relaxation

Emotion control	n	Mean	SD	Difference		CI 95%	P
				Mean	SD		
Before	31	30.03	5.73	7.97	7.08	5.65-10.28	0.001
After	31	38.00	6.05				

4. DISCUSSION

In this study researchers wanted to examine the effect of relaxation techniques to control emotions of patients with tuberculosis. The results in Table 2 confirmed that the value of mean and standard deviation was 30.03 (± 5.72) before the treatment of deep breathing relaxation techniques, and after the treatment the mean and standard deviation was 38.00 (± 6.05) with p value=0.001. The results of this study indicated that deep breathing relaxation techniques significantly effected on controlling emotions of patients with tuberculosis.

This results is supported by the research results conducted by Yilmaz, GS et al (2015) in patients with breast cancer which mentioned that before the intervention group having relaxation exercises, the mean and standard deviation was 42.26 (± 7.49) and after the exercises was 36.20 (± 7.49), with p-value=0.001 [7]. The results are also similar to the study conducted by M.J. Naveen [5] towards patients with Chronic Obstructive Pulmonary Disease (COPD). It was stated that the pretest results against intervention group was 14 (70%) from the patients who had severe anxiety and 6 (30%) from patients who had moderate anxiety. In posttest, afterward, there were only found 2 (10%) patients with mild anxiety and 18 (90%) patients with moderate anxiety [5].

The other similar study was performed by Lin MF et al (2010) towards the undergoing chemotherapy patients with cancer. The study found that before verbal relaxation to the intervention group, the anxiety level reached the mean and standard deviation of 41.07 (13.71) and after verbal relaxation the anxiety level obtained the mean and standard deviation of 31.63 (10.96), with p value 0.001 [8]. This study was also related to the study of Sepasi, N et al [9]. to breast cancer patients. It was stated that the value of mean and standard deviation towards intervention group of relaxation exercises of their emotional well-being pre-test was 9.90 (3.68) and in the post test it obtained 19.40 (2:41) with P value <0.001 [9].

Emotion is a situation that is volatile in the individual that serves as an inner adjustment (adjustment from inside) to the environment to achieve well-being and safety of patients [10]. The psychological impact on tuberculosis patients is depressed, disappointed, angry and anxious. The research conducted by Martin. L.R et al [11] indicated that one of the strongest factors caused non-adherent patients on taking tuberculosis medication is depressed. Depressed patients have non-compliance medication risks by 27%. According to Martin et al [11], patients with tuberculosis will experience cognitive impairment, pessimistic and withdraw from social environment that causes reducing ability to follow treatment regimens.

Emotional control is the tendency to hide and calm the emotions of negative emotions, such as anger, depression or anxiety. Controlling emotions influence the adjustment to living with the disease through much more complex adaptive processes such as strategies for coping with stress [3].

The high capability of controlling emotions was found in female because they are better in keeping interpersonal relations [12]. The ability to control emotions remained stable in adults because at the adult ages the attention and memory aimed at a positive effect by processing a negative stimulus into a positive stimulus [13]. It can be argued that the more age, the more stable the emotional ability. Mahmud in Sobur [10] mentioned some rules for controlling emotions, they are: Deal with the emotions of people, re-interpreted the situation, develop a sense of humor and a realistic attitude. Meanwhile, according to Bast in Hardiyani, the way to control angry emotions when being in unpleasant situation are relaxation, cognitive restructuring and better communication.

5. CONCLUSION

Tuberculosis patients have a tendency to experience emotional disturbance due to the illness. Deep breathing relaxation techniques conducted over three weeks is effective to control emotions of patients with tuberculosis.

ACKNOWLEDGMENTS

The authors would like to thank to the Head and staffs of the Society Pulmonary Health (BKPM) Magelang who has given permission to conduct research, and also to all of research respondents.

REFERENCES

- [1] WHO, "Global Tuberculosis Report [Internet]," vol. 1, 2015. Available from: <https://www.health-e.org.za/wp-content/uploads/2015/10/Global-TB-Report-2015-FINAL-2.pdf>.
- [2] X. B. Paulo and P. Bruno, "Emotional distress in Angolan patients with several types of tuberculosis," *Afr Health Sci.*, vol/issue: 15(2), pp. 2–5, 2015.
- [3] Janowski K., Kurpas D., Kusz J., Mroczek B., Jedynak T., "Emotional control, styles of coping with stress and acceptance of illness among patients suffering from chronic somatic diseases," *Stress Heal*, vol/issue: 30(1), pp. 34–42, 2014. Available from: <http://web.b.ebscohost.com/ehost/pdfviewer/pdfviewer?sid=71430802-36c5-402a-8d4d-6753e2729f56@sessionmgr102&vid=1&hid=106>.
- [4] K. Chalco, *et al.*, "Nurses as providers of emotional support to patients with MDR-TB," pp. 253–61, 2006.
- [5] M N. J., Porkodi A., Akila P., "Psycho-Physiological Parameters Among Patients," *J Sci*, vol/issue: 4(1), pp. 23–6, 2014. Available from: http://www.journalofscience.net/File_Folder/23-26.pdf.
- [6] Dahlan M. S., "Statistik Untuk Kedokteran dan Kesehatan," Jakarta, Salemba Medika, 2013.
- [7] Yilmaz S. G., Arslan S., "Effects of Progressive Relaxation Exercises on Anxiety and Comfort of Turkish Breast Cancer Patients Receiving Chemotherapy," *Asian Pacific J Cancer Prev*, vol. 16, pp. 217–20, 2015. Available from: https://www.researchgate.net/publication/271709071_Effects_of_Progressive_Relaxation_Exercises_on_Anxiety_and_Comfort_of_Turkish_Breast_Cancer_Patients_Receiving_Chemotherapy.
- [8] Lin M., Hsieh Y., Hsu Y., Fetzter S., Hsu M., "Randomised controlled trial of the effect of music therapy and verbal relaxation on chemotherapy-induced anxiety," *J Sci*, issue 8, pp. 988–99, 2011. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21385249>.
- [9] Sepasi N., Qasemi A. A., Narimani M., "A comparison of efficacy of two training techniques of self-management and relaxation on the aspects of quality of life in women with cancer," vol. 6, pp. 18–28, 2014. Available from: <http://bccr.tums.ac.ir/index.php/bccrj/article/view/88>.
- [10] Sobur A., "Psykologi Umum dalam lintasan sejarah," Bandung, Pustaka Setia, 2013.
- [11] Martin L. R., Williams S. L., Haskard K. B., Dimatteo M. R., "The challenge of patient adherence," *Ther Clin Risk Manag*, vol/issue: 1(3), pp. 189–99, 2005. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1661624&tool=pmcentrez&rendertype=abstract>
- [12] D. Coon, J. O. Mitterer, "Psychology a Journey," USA, 2010.
- [13] Nashiro K., Sakaki M., Mather M., "Age differences in brain activity during emotion processing: Reflections of age-related decline or increased emotion regulation?," *Gerontology*, vol/issue: 58(2), pp. 156–63, 2012.

Jurnal 3 - Deep Breathing Relaxation Techniques Improve Emotional Control on Tuberculosis Patients

ORIGINALITY REPORT

13%

SIMILARITY INDEX

8%

INTERNET SOURCES

4%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	doi.org Internet Source	1%
2	Submitted to The Chicago School of Professional Psychology Student Paper	1%
3	Submitted to University of Western Ontario Student Paper	1%
4	umexpert.um.edu.my Internet Source	1%
5	Submitted to University of Ulster Student Paper	1%
6	www.slideshare.net Internet Source	1%
7	digilib.iain-palangkaraya.ac.id Internet Source	1%
8	ebettingsystems.com Internet Source	1%

repository.upi.edu

9	Internet Source	1 %
10	Submitted to Universitas Jenderal Soedirman Student Paper	1 %
11	Submitted to University of Nottingham Student Paper	1 %
12	Noelle M. Chiavetta, Ana R.S. Martins, Inês C.R. Henriques, Felipe Fregni. "Differences in methodological quality between positive and negative published clinical trials", Journal of Advanced Nursing, 2014 Publication	<1 %
13	digilib.uinsgd.ac.id Internet Source	<1 %
14	journals.sagepub.com Internet Source	<1 %
15	A W Harahap, M S Nasution. "Comparison quality of life patients treated with insulin and oral hypoglycemic drugs", IOP Conference Series: Earth and Environmental Science, 2018 Publication	<1 %
16	Ernawati ., Agus Sustiyono, Mulyati .. "The Effectiveness of Ma'tsurat Dhikr in Reducing Blood Sugar Levels in Type 2 Diabetes Mellitus Patients at Dr. Dradjat Prawiranegara	<1 %

Hospital Serang in 2018", KnE Life Sciences, 2019

Publication

-
- | | | |
|----|---|------|
| 17 | repo.unand.ac.id
Internet Source | <1 % |
| 18 | www.frontiersin.org
Internet Source | <1 % |
| 19 | Heather Greenlee, Melissa J. DuPont-Reyes, Lynda G. Balneaves, Linda E. Carlson et al. "Clinical practice guidelines on the evidence-based use of integrative therapies during and after breast cancer treatment", CA: A Cancer Journal for Clinicians, 2017
Publication | <1 % |
| 20 | Adrianus Ola Wuan, Ayorince Herlinalt Gloria Banunu, Norma Tiku Kambuno. "Total Lymfosit Count (TLC) with CD4 in HIV/AIDS Patients at Kupang", Jurnal Teknologi Laboratorium, 2019
Publication | <1 % |
-

Exclude quotes On

Exclude matches Off

Exclude bibliography On