

DAFTAR PUSTAKA

- Alberta Biodiversity Monitoring Program. (2006). *Laboratory protocols for processing Bryophytes (Version 2.0)*. Alberta Biodiversity Monitoring Program.
- Anglana, C., Barozzi, F., Capaci, P., Migoni, D., Rojas, M., Fanizzi, F. P., & Di Sansebastiano, G. P. (2024). Characterization of three species of aquatic mosses in axenic culture for biomonitoring and biotechnological applications. *Aquatic Botany*, 193, Article 103762. <https://doi.org/10.1016/j.aquabot.2024.103762>
- Asakawa, Y., Ludwiczuk, A., & Nagashima, F. (2013). Phytochemical and biological studies of *Bryophytes*. *Phytochemistry*, 91, 52–80. <https://doi.org/10.1016/j.phytochem.2012.04.012>
- Badan Perencanaan Pembangunan Nasional. (2024). *Strategi dan rencana aksi keanekaragaman hayati Indonesia (Indonesian Biodiversity Strategy and Action Plan IBSAP 2025-2045)*. Badan Perencanaan Pembangunan Nasional.
- Bakalin, V., Choi, S. S., & Park, S. J. (2021). Review of *Heteroscyphus Schiffn.*(Lophocoleaceae, Marchantiophyta) in Cambodia. *Journal of Bryology*, 43(1), 52-61. <https://doi.org/10.1080/03736687.2020.1815383>
- Bandyopadhyay, A., & Dey, A. (2022). The ethno-medicinal and pharmaceutical attributes of *Bryophytes*: A review. *Phytomedicine Plus*, 2(2), Article 100255. <https://doi.org/10.1016/j.phyplu.2022.100255>
- Barnabas, S. M. (2016). Keanekaragaman lumut (Bryophyta) di kawasan Danau Tambing Taman Nasional Lore Lindu dan pengembangannya sebagai media pembelajaran. *Mitra Sains*, 4(1), 76–84.
- Benavides, J. C., Duque, A. J., Duivenvoorden, J. F., Vasco, A., Callejas, R., & Stevenson, P. (2011). Effects of forest structure on the diversity of epiphytic *Bryophytes* in Amazonian forests. *Journal of Tropical Ecology*, 27(5), 503–516. <https://doi.org/10.1017/S026646741100021X>
- Benita, A. A., Fitriani, A., Ramadhan, F., Masfurin, F. R., Grandisningtias, G. G., & Fitriyyah, I. (2025). Identifikasi karakter morfologi dan reproduksi pada Thallophyta dan Bryophyta di UIN Sunan Gunung Djati Bandung. *Tumbuhan: Publikasi Ilmu Sosiologi Pertanian dan Ilmu Kehutanan*, 2(1), 82–94.

- Brady, K. U., Kruckeberg, A. R., & Bradshaw Jr, H. D. (2005). Evolutionary ecology of plant adaptation to serpentine soils. *Annual Review of Ecology, Evolution, and Systematics*, 36(1), 243–266. <https://doi.org/10.1146/annurev.ecolsys.36.102403.094325>
- Breil, D. A. (2003). Common and occasional *Bryophytes* of the Virginia Piedmont. *Banisteria*, 21, 3–27.
- Buck, W. R., & Vitt, D. H. (2006). *Pleurocarpous mosses of the world*. Memoirs of the New York Botanical Garden, 98, 1–320.
- Büdel, B., Friedl, T., & Beyschlag, W. (Eds.). (2024). Biology of algae, lichens and *Bryophytes*. *Springer Spektrum*. <https://doi.org/10.1007/978-3-662-65775-1>
- Burdon, F. J., Ramberg, E., Sargac, J., Forio, M. A. E., De Saeyer, N., Mutinova, P. T., ... & McKie, B. G. (2020). Assessing the benefits of forested riparian zones: A qualitative index of riparian integrity is positively associated with ecological status in European streams. *Water*, 12(4), Article 1178. <https://doi.org/10.3390/w12041178>
- Cain, M.L., Bowman, W.D., & Hacker, S.D. (2014). *Ecology (3rd Edition)*. Sinauer Associates.
- Cannone, N., Dalle Fratte, M., Convey, P., Worland, M. R., & Guglielmin, M. (2017). Ecology of moss banks on Signy Island (maritime Antarctic). *Botanical Journal of the Linnean Society*, 184(4), 518–533. <https://doi.org/10.1093/botlinnean/box047>
- Ceter, T., Gözcü, M. C., & Uyar, G. (2018). Spore morphology of some Bartramiaceae species (Lumut) in Turkey. *Communications Faculty of Sciences University of Ankara Series C Biology*, 27(2), 253–262. https://doi.org/10.1501/commuc_0000000222
- Chadijah, A., Sulistiono, S., Affandi, R., Haryani, G. S., & Mashar, M. A. (2022). The reproduction of endemic fish Opudi (*Telmatherina prognatha* Kottelat, 1991) in Lake Matano, South Sulawesi, Indonesia. *International Journal on Advanced Science, Engineering and Information Technology*, 12(2), 697–702. <https://doi.org/10.18517/ijaseit.12.2.14833>
- Crowe, S. A., O'Neill, A. H., Katsev, S., Hehanussa, P., Haffner, G. D., Sundby, B., Mucci, A., & Fowle, D. A. (2008). The biogeochemistry of tropical lakes: A case study from Lake Matano, Indonesia. *Limnology and Oceanography*, 53(1), 319–331. <https://doi.org/10.4319/lo.2008.53.1.0319>

- da Costa, D. P. (1999). Epiphytic bryophyte diversity in primary and secondary lowland rainforests in southeastern Brazil. *The Bryologist*, 102(2), 320–326. <https://doi.org/10.2307/3244261>
- Decker, E. L., & Reski, R. (2020). Mosses in biotechnology. *Current Opinion in Biotechnology*, 61, 21–27. <https://doi.org/10.1016/j.copbio.2019.09.021>
- Dziwak, M., Wróblewska, K., Szumny, A., & Galek, R. (2022). Modern use of bryophytes as a source of secondary metabolites. *Agronomy*, 12(6), Article 1456. <https://doi.org/10.3390/agronomy12061456>
- Eddy, A. (1988). *A handbook of Malesian mosses: Volume 1. Sphagnales to Dicranales*. Natural History Museum Publications.
- Eddy, A. (1990). *A handbook of Malesian mosses: Volume 2. Leucobryaceae to Buxbaumiaceae*. Natural History Museum Publications.
- Ellis, L. T. (2020). Notes on Calymperes subg. Calymperes (Calymperaceae) in the Malesian-Pacific region, with special reference to *C. subulatum*, *C. beccarii* and a new species from Borneo. *Journal of Bryology*, 42(4), 301–315. <https://doi.org/10.1080/03736687.2020.1824880>
- Frahm, J. P. (2003). *Manual of tropical bryology*. Botanisches Institut der Universität.
- Frangedakis, E., Shimamura, M., Villarreal, J. C., Li, F. W., Tomaselli, M., Waller, M., ... & Szövényi, P. (2021). The hornworts: Morphology, evolution and development. *New Phytologist*, 229(2), 735–754. <https://doi.org/10.1111/nph.16874>
- Glime, J. M. (2017). *Bryophyte ecology*. Michigan Technological University.
- Goffinet, B., & Shaw, A. J. (Eds.). (2009). *Bryophyte biology (2nd ed.)*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511754890>
- González, E., Felipe-Lucia, M., Bourgeois, B., Boz, B., Nilsson, C., Palmer, G., & Sher, A. (2017). Integrative conservation of riparian zones. *Biological Conservation*, 211, 20–29. <https://doi.org/10.1016/j.biocon.2016.10.035>
- Gradstein, S. R., & Terken, L. (1981). Studies on Lejeuneaceae subfam. Ptychanthoideae VI. A revision of *Schiffneriolejeunea* sect. *Saccatae* from Asia. *Occasional Papers of the Farlow Herbarium of Cryptogamic Botany*, 71–81. <https://doi.org/10.5962/p.317793>

- Gradstein, S. R. (1992). *The vanishing tropical rain forest as an environment for Bryophytes and lichens*. In J. W. Bates & A. M. Farmer (Eds.), *Bryophytes and lichens in a changing environment* (pp. 232–256). Clarendon Press. <https://doi.org/10.1093/oso/9780198542919.003.0009>
- Gradstein, S. R. (2011). *Guide to the liverworts and hornworts of Java*. Seameo Biotrop.
- Gradstein, S. R. (2020). The genera of Lejeuneaceae (Marchantiophyta) of tropical America – an update. *Nova Hedwigia, Beihefte*, 150, 81–96. <https://doi.org/10.1127/nova-suppl/2020/081>
- Gradstein, S. R., & León-Yáñez, S. (2020). Liverwort diversity in Polylepis pautas forests of Ecuador under different climatic conditions. *Neotropical Biodiversity*, 6(1), 138–146. <https://doi.org/10.1080/23766808.2020.1802928>
- Gradstein, S. R., Churchill, S. P., & Salazar-Allen, N. (2001). *Guide to the Bryophytes of tropical America*. Memoirs of the New York Botanical Garden, 86, 1–577.
- Gu, Z., Gu, L., Eils, R., Schlesner, M., & Brors, B. (2014). circlize implements and enhances circular visualization in R. *Bioinformatics*, 30(19), 2811–2812. <https://doi.org/10.1093/bioinformatics/btu393>
- Haffner, G. D., Hehanussa, P. E., & Hartoto, D. (2001). *The biology and physical processes of large lakes of Indonesia: Lakes Matano and Towuti*. In M. Munawar (Ed.), *The Great Lakes of the World: Food-web, health and integrity* (pp. 183–192). Backhuys Publishers.
- Hassan, F. M., Al-Zubaidi, N. A. J., El-Sheekh, M. M., Youssef, O. S., Salman, J. M., Douabul, A., & Panhwar, S. K. (2023). Temporal and spatial distribution of epiphyton with diversity indices in a lotic ecosystem. *Egyptian Journal of Aquatic Biology and Fisheries*, 27(6), 815–836. <https://doi.org/10.21608/ejabf.2023.328312>
- He, X., Sun, Y., & Zhu, R. L. (2013). The oil bodies of liverworts: Unique and important organelles in land plants. *Critical Reviews in Plant Sciences*, 32(5), 293–302. <https://doi.org/10.1080/07352689.2013.765775>
- Heinrichs, J., Kettunen, E., Lee, G. E., Marzaro, G., Pócs, T., Ragazzi, E., ... & Schmidt, A. R. (2015). Lejeuneaceae (Marchantiophyta) from a species-rich taphocenosis in Miocene Mexican amber, with a review of liverworts fossilised in amber. *Review of Palaeobotany and Palynology*, 221, 59–70. <https://doi.org/10.1016/j.revpalbo.2015.06.002>

- Heinrich, A. (2023). Bryophytes of Santa Barbara. <https://bryophytessb.com/liverworts/porella-bolanderi-9/>. (Diakses pada 13 Juni 2026).
- Herder, F., Schliewen, U. K., Geiger, M. F., Hadiaty, R. K., Gray, S. M., McKinnon, J. S., ... & Pfaender, J. (2012). Alien invasion in Wallace's Dreamponds: records of the hybridogenic "flowerhorn" cichlid in Lake Matano, with an annotated checklist of fish species introduced to the Malili Lakes system in Sulawesi. *Aquatic Invasions*, 7(4). <http://dx.doi.org/10.3391/ai.2012.7.4.009>
- Holt, E. A., & Miller, S. W. (2011). Bioindicators: Using organisms to measure. *Nature*, 3, 8-13.
- Horn, A., Pascal, A., Lončarević, I., Volpato Marques, R., Lu, Y., Miguel, S., ... & Simonsen, H. T. (2021). Natural products from *Bryophytes*: From basic biology to biotechnological applications. *Critical Reviews in Plant Sciences*, 40(3), 191–217. <https://doi.org/10.1080/07352689.2021.1922013>
- Huttunen, S., & Quandt, D. (2006). *Phylogenetic relationships within the moss family Meteoriaceae in the light of different datasets, alignment and analysis methods*. In A. E. Newton & R. S. Tangney (Eds.), *Pleurocarpous Mosses: Systematics and Evolution* (pp. 141–157). CRC Press. <https://doi.org/10.1201/9781420006997.ch7>
- Huttunen, S., Bell, N., & Hedenäs, L. (2018). The evolutionary diversity of mosses—taxonomic heterogeneity and its ecological drivers. *Critical Reviews in Plant Sciences*, 37(2-3), 128–174. <https://doi.org/10.1080/07352689.2018.1482424>
- Hylander, K., Dynesius, M., Jonsson, B. G., & Nilsson, C. (2005). Substrate form determines the fate of *Bryophytes* in riparian buffer strips. *Ecological Applications*, 15(2), 674–688. <https://doi.org/10.1890/04-0570>
- Jaccard, P. (1901). Étude comparative de la distribution florale dans une portion des Alpes et des Jura. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 37, 547–579.
- Jiang, T., Yang, X., Zhong, Y., Tang, Q., Liu, Y., & Su, Z. (2018). Species composition and diversity of ground bryophytes across a forest edge-to-interior gradient. *Scientific Reports*, 8(1), Article 11868. <https://doi.org/10.1038/s41598-018-30281-w>
- Jumari, & Soeprbowati, T. R. (2024). Vegetasi riparian Telaga Pengilon dan gangguan antropogenik. *Jurnal Ilmu Lingkungan*, 22(2), 455–463. <https://doi.org/10.14710/jil.22.2.455-463>

- Johansson, L., & Renman, G. (1994). The use of aquatic bryophytes as bioindicators. *Internationale Vereinigung für theoretische und angewandte Limnologie: Verhandlungen*, 25(3), 2026-2028. <https://doi.org/10.1080/03680770.1992.11900552>
- Kent, M. (2012). *Vegetation description and data analysis: A practical approach* (2nd ed.). John Wiley & Sons.
- Koponen, T., Ignatova, E. A., Kuznetsova, O. I., & Ignatov, M. S. (2012). Philonotis (Bartramiaceae, Lumut) in Russia. *Arctoa*, 21, 21-62.
- Koroleva, Y., Napreenko, M., Baymuratov, R., & Schefer, R. (2020). Bryophytes as a bioindicator for atmospheric deposition in different coastal habitats (a case study in the Russian sector of the Curonian Spit, South-Eastern Baltic). *International Journal of Environmental Studies*. <https://doi.org/10.1080/00207233.2019.1594301>
- Lathifah, A. N., Guo, Y., Sakagami, N., Suda, W., Higuchi, M., Nishizawa, T., ... & Ohta, H. (2019). Comparative characterization of bacterial communities in moss-covered and unvegetated volcanic deposits of Mount Merapi, Indonesia. *Microbes and environments*, 34(3), 268-277. <https://doi.org/10.1264/jsme2.ME19041>
- Lee, G. E. (2013). A systematic revision of the genus *Lejeunea* Lib. (Marchantiophyta: Lejeuneaceae) in Malaysia. *Cryptogamie, Bryologie*, 34(4), 381-484. <https://doi.org/10.7872/cryb.v34.iss4.2013.381>
- Lee, G. E., & Gradstein, S. R. (2021). *Guide to the genera of liverworts and hornworts of Malaysia*. Hattori Botanical Laboratory.
- Legendre, P., & Legendre, L. (2012). *Numerical ecology* (3rd ed.). Elsevier.
- Lehmusluoto, P., Machbub, B., Terangna, N., Achmad, F., Boer, L., Brahmana, S. S., Priadi, B., Setiadji, B., Sayuman, O., & Margana, A. (1995). *National inventory of the major lakes and reservoirs in Indonesia*. General Limnology: FAO/FINNIDA.
- Ligrone, R., Duckett, J. G., & Renzaglia, K. S. (2012). Major transitions in the evolution of early land plants: A bryological perspective. *Annals of Botany*, 109(5), 851-871. <https://doi.org/10.1093/aob/mcs017>
- Lin, W., Wang, Y., Coudert, Y., & Kierzkowski, D. (2021). Leaf morphogenesis: Insights from the moss *Physcomitrium patens*. *Frontiers in Plant Science*, 12, Article 736212. <https://doi.org/10.3389/fpls.2021.736212>

- Magurran, A. E. (2004). *Measuring biological diversity*. Blackwell Publishing.
- Manju, C. N., Martin, K. P., Sreekumar, V. B., & Rajesh, K. P. (2012). Morphological and molecular differentiation of *Aerobryopsis eravikulamensis* and closely related taxa of the Western Ghats of India. *The Bryologist*, 115(1), 42–50. <https://doi.org/10.1639/0007-2745-115.1.42>
- McCune, B., & Grace, J. B. (2002). *Analysis of ecological communities*. MjM Software Design.
- Modi, G., Arora, B., Bhardwaj, N., & Babita. (2023). An initiatory review on bryophytes diversity. *International Journal for Research in Applied Science and Engineering Technology*, 11(2). <https://doi.org/10.22214/ijraset.2023.49013>
- Mohamed, H., & Reese, W. D. (1985). Syrrhopodon (Musci: Calymperaceae) in Malaysia and adjacent regions. *Bryologist*, 88(3). 223-254. <https://doi.org/10.2307/3243033>
- Morin, P.J. (2011). *Community Ecology (2nd Edition)*. Wiley-Blackwell.
- Muamar, A., Anggraito, Y. U., & Retnoningsih, A. (2021, June). The diversity of moss in the Cemoro Kandang hiking trail, Mount Lawu and the Baturraden botanical gardens, Central Java. In *Journal of Physics: Conference Series* (Vol. 1918, No. 5, p. 052039). IOP Publishing. <https://doi.org/10.1088/1742-6596/1918/5/052039>
- Naren, G. W., & Jia, Y. (2020). Taxonomic study on Meteoriaceae I. Reappraisal of some species of *Barbella* and *Pseudotrachypus*. *Journal of Bryology*, 42(4), 333-342. <https://doi.org/10.1080/03736687.2020.1839247>
- Nóbrega, R., Ziembowicz, T., Torres, G., Guzha, A., Amorim, R., Cardoso, D., Johnson, M., Santos, T., Couto, E., & Gerold, G. (2019). Ecosystem services of a functionally diverse riparian zone in the Amazon–Cerrado agricultural frontier. *Global Ecology and Conservation*. <https://doi.org/10.32942/osf.io/85ypg>.
- Oishi, Y. (2018). Evaluation of the water-storage capacity of *Bryophytes* along an altitudinal gradient from temperate forests to the alpine zone. *Forests*, 9(7), 433. <https://doi.org/10.3390/f9070433>
- Pasaribu, N. (2013). Studi pendahuluan lumut di Lau Kawar, Kabupaten Karo. *Prosiding SEMIRATA 2013*, 1(1), 45-51.

- Passarella, M. D. A., & Luizi-Ponzo, A. P. (2019). Palynology of *Amphidium Schimp.* (Amphidiaceae M. Stech): Can spore morphology circumscribe the genus?. *Acta Botanica Brasilica*, 33(1), 135–140. <https://doi.org/10.1590/0102-33062018abb0235>
- Patra, I. K., & Hayat, T. N. P. S. (2019). Prospek Pengembangan Pariwisata Danau Matano Dalam Meningkatkan Pendapatan Asli Daerah (Pad) Kabupaten Luwu Timur. *Equilibrium: Jurnal Ilmiah Ekonomi, Manajemen dan Akuntansi*, 7(2). <https://doi.org/10.35906/je001.v7i2.324>
- Podengge, E., & Puspitasari, D. J. (2023). Aktivitas penduduk terhadap pencemaran air Danau Matano di Desa Sorowako Kecamatan Nuha Kabupaten Luwu Timur. *Jurnal Gawalise*, 2(1), 11–20.
- Printarakul, N., & Meeinkuirt, W. (2022). The bryophyte community as bioindicator of heavy metals in a waterfall outflow. *Scientific reports*, 12(1), 6942. <https://doi.org/10.1038/s41598-022-10980-9>
- Pursell, R. A. (2007). *Fissidentaceae. Flora Neotropica Monograph 101*. New York Botanical Garden Press.
- Putrika, A., Salamah, A., Atria, M., Hendrayanti, D., Winarni, N. L., Purba, E. P., ... & Ariyanti, N. S. (2023). Bryophyte diversity and atmospheric pollution in a residential area and an industrial urban forest in Jakarta, Indonesia. *Biotropia*, 30(3), 355–364. <https://doi.org/10.11598/btb.2023.30.3.1960>
- Putrika, A., Salamah, A., Winarni, N. L., Hendrayanti, D., & Dewi, C. (2026). The epiphytic bryophyte diversity across urban land-use type in Southern Jakarta. *Journal of Urban Ecology*, 12(1), juag002. <https://doi.org/10.1093/jue/juag002>
- Qiao, Z., Lin, X., Zhang, T., Luo, S., Wang, C., Zhou, W., & Luo, L. (2025). A review of research progress on the application of bryophytes in the ecological restoration of mining areas of China. *Frontiers in Environmental Science*, 13, 1610193. <https://doi.org/10.3389/fenvs.2025.1610193>
- Reese, W. D., Mohamed, H., & Mohamed, A. D. (1986). A synopsis of *Mitthyridium* (Musci: Calymperaceae) in Malaysia and adjacent regions. *Bryologist*, 49–58. <https://doi.org/10.2307/3243077>
- Rohlf, F. J. (1970). Adaptive hierarchical clustering schemes. *Systematic Zoology*, 19(1), 58–82. <https://doi.org/10.2307/2412026>
- Salazar Allen, N. (1993). A revision of the pantropical moss genus *Leucophanes* Brid. *Bryophytorum Bibliotheca*, 46, 1–281.

- Samarakkody, S. P., Ruklani, N. C. S., & Rubasinghe, S. C. K. (2018). Three new species records of leafy liverworts (Marchantiophyta, Jungermanniidae) to Sri Lanka. *Lindbergia*, 41(1), Article 01110. <https://doi.org/10.25227/linbg.01110>
- Sen, K., Khan, M. I., Paul, R., Ghoshal, U., & Asakawa, Y. (2023). Recent advances in the phytochemistry of Bryophytes: Distribution, structures and biological activity of Bibenzyl and Bisbibenzyl compounds. *Plants*, 12(24), 4173. <https://doi.org/10.3390/plants12244173>
- Seppelt, R. D., Meagher, D., Cairns, A., & Franks, A. (2021). The moss family Calymperaceae (Lumut) in Australia Part 1: Introduction and key to genera. *Telopea*, 24, 247–252. <https://doi.org/10.7751/telepeal5012>
- Shen, T., Corlett, R. T., Collart, F., Kasprzyk, T., Guo, X. L., Patiño, J., ... & Vanderpoorten, A. (2022). Microclimatic variation in tropical canopies: A glimpse into the processes of community assembly in epiphytic bryophyte communities. *Journal of Ecology*, 110(12), 3023–3038. <https://doi.org/10.1111/1365-2745.13978>
- Shevock, J. R., Ma, W. Z., & Akiyama, H. (2017). Diversity of the rheophytic condition in *Bryophytes*: Field observations from multiple continents. *Bryophyte Diversity and Evolution*, 39(1), 75–93. <https://doi.org/10.11646/bde.39.1.12>
- Sim-Sim, M., Afonina, O. M., Almeida, T., Désamuré, A., Laenen, B., Garcia, C. A., ... & Stech, M. (2017). Integrative taxonomy reveals too extensive lumping and a new species in the moss genus *Amphidium* (Bryophyta). *Systematics and biodiversity*, 15(5), 451–463. <https://doi.org/10.1080/14772000.2016.1271059>
- Singh, R., Tiwari, A. K., & Singh, G. S. (2021). Managing riparian zones for river health improvement: an integrated approach. *Landscape and ecological engineering*, 17(2), 195–223. <https://doi.org/10.1007/s11355-020-00436-5>
- Singh, S., & Srivastava, K. (2013). Lumut as green brain: Unique and indispensable small creature. *International Journal of Pharmaceutical Sciences Review and Research*, 23(2), 28–35.
- Siwach, A., Kaushal, S., & Baishya, R. (2021). Effect of Mosses on physical and chemical properties of soil in temperate forests of Garhwal Himalayas. *Journal of Tropical Ecology*, 37(3), 126–135. <https://doi.org/10.1017/S0266467421000249>

- Slate, M. L., Antoninka, A., Bailey, L., Berdugo, M. B., Callaghan, D. A., Cárdenas, M., ... & Coe, K. K. (2024). Impact of changing climate on bryophyte contributions to terrestrial water, carbon, and nitrogen cycles. *New Phytologist*, 242(6), 2411–2429. <https://doi.org/10.1111/nph.19741>
- Sorenson, T. (1948). A method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on Danish commons. *Biol Skar*, 5, 1.
- Spitale, D. (2016). The interaction between elevational gradient and substratum reveals how *Bryophytes* respond to the climate. *Journal of Vegetation Science*, 27(4), 844–853. <https://doi.org/10.1111/jvs.12401>
- Sporn, S. G., Bos, M. M., Hoffstätter-Müncheberg, M., Kessler, M., & Gradstein, S. R. (2009). Microclimate determines community composition but not richness of epiphytic understory bryophytes of rainforest and cacao agroforests in Indonesia. *Functional Plant Biology*, 36(2), 171–179. <https://doi.org/10.1071/FP08197>
- Su, X., Qi, T., Li, Y., Wang, W., Li, D., Yang, X., & Hao, J. (2025). Bryophyte Community Composition and Diversity as Bioindicators of Elevational Zonation in Tropical Rainforests in Hainan Island, China. *Plants*, 14(20), 3209. <https://doi.org/10.3390/plants14203209>
- Sukamto, D. S. (2023). Identifikasi Keanekaragaman Lumut Hati di Danau Ranupani Lumajang. *BIO-CONS: Jurnal Biologi Dan Konservasi*, 5(1), 55–61. <https://doi.org/10.31537/biocons.v5i1.1100>
- Sukkharak, P. (2015). A systematic monograph of the genus *Thysananthus* (Lejeuneaceae, Marchantiophyta). *Phytotaxa*, 193(1), 1–81. <https://doi.org/10.11646/phytotaxa.193.1.1>
- Sulastri, S., Harsono, E., Ridwansyah, I., Dina, R., & Melati, I. (2020). Endemic and invasive species of Lake Matano and allowable suspended solid load to sustain high species endemism. *IOP Conference Series: Earth and Environmental Science*, 535(1), Article 012026. <https://doi.org/10.1088/1755-1315/535/1/012026>
- Turetsky, M. R. (2003). The role of *Bryophytes* in carbon and nitrogen cycling. *The Bryologist*, 106(3), 395–409. [https://doi.org/10.1639/0007-2745\(2003\)106\[0395:trobic\]2.0.co;2](https://doi.org/10.1639/0007-2745(2003)106[0395:trobic]2.0.co;2)
- Turetsky, M. R., Bond-Lamberty, B., Euskirchen, E., Talbot, J., Frohling, S., McGuire, A. D., & Tuittila, E. S. (2012). The resilience and functional role of moss in boreal and arctic ecosystems. *New Phytologist*, 196(1), 49–67. <https://doi.org/10.1111/j.1469-8137.2012.04222.x>

- Utomo, A. A., Khotimperwati, L., & Jumari. (2022). Lejeuneaceae (Marchantiophyta) in Curug Semirang, Central Java, Indonesia. *Biogenesis: Jurnal Ilmiah Biologi*, 10(2), 168–180. <https://doi.org/10.24252/bio.v10i2.29903>
- Vaillant, J. J., Haffner, G. D., & Cristescu, M. E. (2011). The ancient lakes of Indonesia: towards integrated research on speciation. *Integrative and Comparative Biology*, 51(4), 634–643. <https://doi.org/10.1093/icb/ucr101>
- van der Maarel, E. & Franklin, J. (2013). *Vegetation Ecology (2nd Edition)*. Wiley-Blackwell.
- van Steenis, C. G. G. J. (1981). *Rheophytes of the world*. Sijthoff & Noordhoff. <https://doi.org/10.1007/978-94-009-8121-8>
- Vanderpoorten, A., & Goffinet, B. (2009). *Introduction to Bryophytes*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511754890>
- Vega, M., Mazón, M., Sarango-Tandazo, C., Álvaro-Alba, W., & Benítez, Á. (2026). Land use effects on epiphytic bryophyte diversity in riparian Andean ecosystems of southern Ecuador. *Urban Ecosystems*, 29(2), Article 88. <https://doi.org/10.1007/s11252-025-01642-x>
- von Rintelen, K., & Cai, Y. (2009). Radiation of endemic species flocks in ancient lakes: Systematic revision of the freshwater shrimp Caridina H. Milne Edwards, 1837 (Crustacea: Decapoda: Atyidae) from the ancient lakes of Sulawesi, Indonesia, with the description of eight new species. *The Raffles Bulletin of Zoology*, 57(2), 343–452.
- Wierzcholska, S., Dyderski, M. K., Pielech, R., Gazda, A., Smoczyk, M., Malicki, M., ... & Jagodziński, A. M. (2018). Natural forest remnants as refugia for bryophyte diversity in a transformed mountain river valley landscape. *Science of The Total Environment*, 640, 954–964. <https://doi.org/10.1016/j.scitotenv.2018.05.333>
- Wolski, G. J., Sadowska, B., Fol, M., Podsędek, A., Kajszyk, D., & Kobylińska, A. (2021). Cytotoxicity, antimicrobial and antioxidant activities of mosses obtained from open habitats. *PLoS One*, 16(9), e0257479. <https://doi.org/10.1371/journal.pone.0257479>
- Wulf, P., & Pearson, R. (2017). Mossy stones gather more bugs: Moss as habitat, nurseries and refugia for tropical stream invertebrates. *Hydrobiologia*, 790, 167–182. <https://doi.org/10.1007/s10750-016-3028-8>

- Yuzawa, Y. I., Mues, R., & Hattori, S. (1987). Morphological and chemical studies on the taxonomy of 14 *Frullania* species, subgenus *Chonantheria*. *The Journal of the Hattori Botanical Laboratory*, 63, 425–436.
- Zander RH. 1993. Genera of Pottiaceae: mosses of harsh environments. *Bulletin of the Buffalo Society of Natural Sciences*, 32, 1–378.
- Zhang, J., Fu, X. X., Li, R. Q., Zhao, X., Liu, Y., Li, M. H., ... & Chen, Z. D. (2020). The hornwort genome and early land plant evolution. *Nature Plants*, 6(2), 107–118. <https://doi.org/10.1038/s41477-020-0618-2>
- Zheng, T. X. (2024). First Record of *Heteroscyphus planus* (Lophocoleaceae) from India, with notes on the liverwort collections from India in the Herbarium of the Hattori Botanical Laboratory (NICH). *Acta Phytotaxonomica et Geobotanica*, 75(3), 185-189. <https://doi.org/10.18942/apg.202412>
- Zheng, G., Gu, J., Zhao, W., Zhang, Y., Guan, Z., Lei, M., & He, C. (2024). Spatial, geographical, climatic, and edaphic influences on moss community structure: A case study from Qinhuangdao, China. *Forests*, 15(3), 424. <https://doi.org/10.3390/f15030424>
- Zotz, G., & Bader, M. Y. (2009). *Epiphytic plants in a changing world-global: change effects on vascular and non-vascular epiphytes*. In *Progress in botany* (pp. 147-170). Berlin, Heidelberg: Springer Berlin Heidelberg.